

**THE NUMBER OF FAMILIES OF SOLUTIONS
OF DECOMPOSABLE FORM EQUATIONS**

J.-H. EVERTSE (LEIDEN) AND K. GYÖRY (DEBRECEN) ^{*)}

1. INTRODUCTION

In [16], Schmidt introduced the notion of family of solutions of norm form equations and showed that there are only finitely many such families. In [18], Voutier gave an explicit upper bound for the number of families. Independently, in [5], Györy extended the notion of family of solutions of norm form equations to decomposable form equations and gave an explicit upper bound for the number of families. In this paper, we obtain a significant improvement of the upper bounds of Voutier and Györy, by applying the results from Evertse [4].

Let β be a non-zero rational integer. Further, let M denote an algebraic number field of degree r and $l(\mathbf{X}) = \alpha_1 X_1 + \cdots + \alpha_m X_m$ a linear form with coefficients in M . There is a non-zero $c \in \mathbf{Q}$ such that the norm form

$$(1.1) \quad F(\mathbf{X}) = cN_{M/\mathbf{Q}}(l(\mathbf{X})) = c \prod_{i=1}^r (\alpha_1^{(i)} X_1 + \cdots + \alpha_m^{(i)} X_m)$$

has its coefficients in \mathbf{Z} . Here, we denote by $\alpha^{(1)}, \dots, \alpha^{(r)}$ the conjugates of $\alpha \in M$. We deal among other things with *norm form equations* of the shape

$$F(\mathbf{x}) = \pm\beta \text{ in } \mathbf{x} \in \mathbf{Z}^m.$$

It is more convenient for us to consider the equivalent equation which is also called a norm form equation,

$$(1.2) \quad cN_{M/\mathbf{Q}}(x) = \pm\beta \text{ in } x \in \mathcal{M},$$

where \mathcal{M} is the \mathbf{Z} -module $\{x = l(\mathbf{x}) : \mathbf{x} \in \mathbf{Z}^m\}$ which is contained in M .

In 1971, Schmidt [15] proved his fundamental result, that (1.2) has only finitely many solutions if \mathcal{M} satisfies some natural non-degeneracy condition. Later, Schmidt

^{*)}Research was supported in part by Grants 16975 and 16791 from the Hungarian National Foundation for Scientific Research and by the Foundation for Hungarian Higher Education and Research.

[16] dealt also with the case that \mathcal{M} is degenerate and showed that in that case, the set of solutions of (1.2) can be divided in a natural way into families, and is the union of finitely many such families. Below, we give a precise definition of a family of solutions of (1.2); here we mention that it is a coset $x\mathfrak{U}_{\mathcal{M},J}$ contained in \mathcal{M} , where x is a solution of (1.2) and $\mathfrak{U}_{\mathcal{M},J}$ is a particular subgroup of finite index in the unit group of the ring of integers of some subfield J of M . Schmidt's results have been generalised to equations of the type

$$(1.3) \quad cN_{M/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M} ,$$

where K is an algebraic number field, \mathcal{O}_S is the ring of S -integers in K for some finite set of places S , \mathcal{O}_S^* is the unit group of \mathcal{O}_S , c, β are elements of $K^* = K \setminus \{0\}$, M is a finite extension of K , and \mathcal{M} is a finitely generated \mathcal{O}_S -module contained in M . In fact, Schlickewei [13] proved the analogue of Schmidt's result on families of solutions in case that \mathcal{O}_S is contained in \mathbf{Q} , and Laurent [9] generalised this to arbitrary algebraic number fields K . The main tools in the proofs of these results were Schmidt's Subspace theorem and Schlickewei's generalisation to the p -adic case and to number fields.

In [5], Gyóry generalised the concept of family of solutions to *decomposable form equations* over \mathcal{O}_S , i.e. to equations of the form

$$(1.4) \quad F(\mathbf{x}) \in \beta\mathcal{O}_S^* \text{ in } \mathbf{x} = (x_1, \dots, x_m) \in \mathcal{O}_S^m ,$$

where K, S are as above, β is a non-zero element of \mathcal{O}_S and $F(\mathbf{X}) = F(X_1, \dots, X_m)$ is a decomposable form with coefficients in \mathcal{O}_S , that is, F can be expressed as a product of linear forms in m variables with coefficients in some extension of K . We can reformulate (1.4) in a shape similar to (1.3) as follows. According to [1], pp. 77-81, there are finite extension fields M_1, \dots, M_t of K , linear forms $l_j(\mathbf{X}) = \alpha_{1j}X_1 + \dots + \alpha_{mj}X_m$ with coefficients in M_j for $j = 1, \dots, t$ and $c \in K^*$ such that

$$(1.5) \quad F(\mathbf{X}) = c \prod_{j=1}^t N_{M_j/K}(l_j(\mathbf{X})) .$$

Now let

$$A = M_1 \oplus \dots \oplus M_t$$

be the direct K -algebra sum of M_1, \dots, M_t , that is, the cartesian product $M_1 \times \dots \times M_t$ endowed with coordinatewise addition and multiplication. If we express an element of A as $(\alpha_1, \dots, \alpha_t)$, then we implicitly assume that $\alpha_j \in M_j$ for $j = 1, \dots, t$. We define the *norm* $N_{A/K}(a)$ of $a = (\alpha_1, \dots, \alpha_t) \in A$ to be the determinant of the K -linear map $x \mapsto ax$ from A to itself. This norm is known to be multiplicative. Further, we have

$$(1.6) \quad N_{A/K}(a) = N_{M_1/K}(\alpha_1) \cdots N_{M_t/K}(\alpha_t)$$

where $N_{M_j/K}$ is the usual field norm. Note that the \mathcal{O}_S -module

$$\mathcal{M} = \{x = (l_1(\mathbf{x}), \dots, l_t(\mathbf{x})) : \mathbf{x} \in \mathcal{O}_S^m\}$$

is contained in A . Now (1.5) and (1.6) imply that eq. (1.4) is equivalent to

$$(1.7) \quad cN_{A/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M} ;$$

(1.7) will also be referred to as a decomposable form equation. In [5], Győry showed that the set of solutions of (1.7) is the union of finitely many families. Further, in [5] he extended some of his results to decomposable form equations over arbitrary finitely generated integral domains over \mathbf{Z} .

In [17], Schmidt made a further significant advancement by deriving, as a consequence of his quantitative Subspace theorem, an explicit upper bound for the number of solutions of norm form equation (1.2) over \mathbf{Z} for every non-degenerate module \mathcal{M} . Schlickewei proved a p -adic generalisation of Schmidt's quantitative Subspace theorem and used this to derive an explicit upper bound for the number of solutions of S -unit equations [14]. Among others, this was used by Győry [5] to obtain an explicit upper bound for the number of families of solutions of decomposable form equation (1.7). Independently, Voutier [18] obtained upper bounds similar to Győry's for the number of families of solutions of norm form equation (1.3), in the special case that $K = \mathbf{Q}$, $\beta = 1$. Recently, Evertse [4] improved the results of Schmidt and Schlickewei just mentioned. In this paper, we apply the results from [4] to obtain an upper bound for the number of families of solutions of (1.7) which is much sharper than Győry's and Voutier's (cf. Theorem 1 in Section 1.2).

In Section 1.1 we introduce the necessary terminology. In Section 1.2 we state our main results (Theorems 1 and 2) and some corollaries. In particular, in Corollary 2 we give an upper bound for the number of \mathcal{O}_S^* -cosets of solutions of (1.7) in case that that number is finite; here, an \mathcal{O}_S^* -coset is a set $x\mathcal{O}_S^* = \{\varepsilon x : \varepsilon \in \mathcal{O}_S^*\}$ where x is a fixed solution of (1.7). Further, in Section 2 we derive from Theorem 1 an asymptotic formula (cf. Corollary 4) for the number of \mathcal{O}_S^* -cosets of solutions of (1.7), in case that this number is infinite. The other sections are devoted to the proofs of Theorems 1 and 2.

1.1. Terminology.

Here and in the sequel we use the following notation: the unit group of a ring R with 1 is denoted by R^* and for $x \in R$ and a subset H of R we define $xH := \{xh : h \in H\}$. Let K be an algebraic number field. Denote by \mathcal{O}_K the ring of integers and by M_K the collection of places (equivalence classes of absolute values) on K . Recall that M_K consists of finitely many infinite (i.e. archimedean) places (the number of these being $r_1 + r_2$ where r_1, r_2 denote the number of isomorphic embeddings of K into \mathbf{R} and the number of complex conjugate pairs of isomorphic embeddings of K into \mathbf{C} , respectively) and of infinitely many finite (non-archimedean) places which may be identified with the prime ideals of \mathcal{O}_K . For every $v \in M_K$ we choose an absolute value $|\cdot|_v$ from v . Now let S be a finite subset of M_K containing all infinite places. The ring of S -integers and its unit group, the group of S -units, are defined by

$$\mathcal{O}_S = \{x \in K : |x|_v \leq 1 \text{ for } v \notin S\}, \quad \mathcal{O}_S^* = \{x \in K : |x|_v = 1 \text{ for } v \notin S\},$$

respectively, where ' $v \notin S$ ' means ' $v \in M_K \setminus S$.' For a finite extension J of K , we denote by $\mathcal{O}_{J,S}$ the integral closure of \mathcal{O}_S in J .

We first introduce families of solutions for norm form equations

$$(1.3) \quad cN_{M/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M},$$

where, as before, M is a finite extension of K , \mathcal{M} is a finitely generated \mathcal{O}_S -module contained in M and c, β are elements of K^* . Let $V := K\mathcal{M}$ be the K -vector space generated by \mathcal{M} . For a subfield J of M containing K , define the sets

$$(1.8) \quad V^J = \{x \in V : xJ \subseteq V\}, \quad \mathcal{M}^J = V^J \cap \mathcal{M}.$$

As is easily seen, we have $\lambda x \in V^J$ for $x \in V^J$, $\lambda \in J$. Further, define the subgroup of the unit group of $\mathcal{O}_{J,S}$,

$$(1.9) \quad \mathfrak{U}_{\mathcal{M},J} := \{\varepsilon \in \mathcal{O}_{J,S}^* : \varepsilon\mathcal{M}^J = \mathcal{M}^J\}.$$

For instance from Lemma 9 of [5] it follows that $\mathfrak{U}_{\mathcal{M},J}$ has finite index in $\mathcal{O}_{J,S}^*$. Note that $N_{M/K}(\varepsilon) \in \mathcal{O}_S^*$ for $\varepsilon \in \mathfrak{U}_{\mathcal{M},J}$. Hence if $x \in \mathcal{M}^J$ is a solution of (1.3) then so is every element of the coset $x\mathfrak{U}_{\mathcal{M},J}$. Such a coset is called a *family of solutions* (or rather an *(\mathcal{M}, J) -family of solutions*) of (1.3). Laurent [9] proved the generalisation of Schmidt's result that the set of solutions of eq. (1.3) is the union of at most finitely many families.

Now let $A = M_1 \oplus \cdots \oplus M_t$ be the direct K -algebra sum of finite extension fields M_1, \dots, M_t of K . Note that A has unit element $1_A = (1, \dots, 1)$ (t times) where 1 is the unit element of K and that the unit group of A is $A^* = \{(\xi_1, \dots, \xi_t) \in A : \xi_1 \cdots \xi_t \neq 0\}$. For each K -subalgebra B of A , denote by $\mathcal{O}_{B,S}$ the integral closure of \mathcal{O}_S in B . Thus,

$$\mathcal{O}_{A,S} = \mathcal{O}_{M_1,S} \oplus \cdots \oplus \mathcal{O}_{M_t,S}$$

is the direct sum of the integral closures of \mathcal{O}_S in M_1, \dots, M_t , respectively, and

$$\mathcal{O}_{B,S} = \mathcal{O}_{A,S} \cap B$$

for each K -subalgebra B of A . From these facts and (1.6) it follows easily, that for $b \in \mathcal{O}_{A,S}$ we have $N_{A/K}(b) \in \mathcal{O}_S$ and that for b in the unit group $\mathcal{O}_{A,S}^*$ we have $N_{A/K}(b) \in \mathcal{O}_S^*$.

Let $c, \beta \in K^*$, let \mathcal{M} be a finitely generated \mathcal{O}_S -module contained in A , and consider the equation

$$(1.7) \quad cN_{A/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M}.$$

Families of solutions of (1.7) are defined in precisely the same way as for (1.3), but now the role of the subfields J of M in (1.3) is played by the K -subalgebras B of A that contain the unit element 1_A of A . Thus, let $V := K\mathcal{M}$ be the K -vector space, contained in A , generated by \mathcal{M} and for each K -subalgebra B of A with $1_A \in B$ define the sets

$$(1.10) \quad V^B := \{x \in V : xB \subseteq V\}, \quad \mathcal{M}^B := V^B \cap \mathcal{M}$$

and the subgroup of the unit group of $\mathcal{O}_{B,S}$,

$$(1.11) \quad \mathfrak{U}_{\mathcal{M},B} := \{\varepsilon \in \mathcal{O}_{B,S}^* : \varepsilon\mathcal{M}^B = \mathcal{M}^B\}$$

which is known to have finite index $[\mathcal{O}_{B,S}^* : \mathfrak{U}_{\mathcal{M},B}]$ in $\mathcal{O}_{B,S}^*$ (cf. [5], Lemma 9). Clearly, V^B is closed under multiplication by elements of B (and in fact the largest subspace of V with this property). A(n (\mathcal{M}, B) -) *family of solutions* of (1.7) is a coset $x\mathfrak{U}_{\mathcal{M},B}$, where B is a K -subalgebra of A containing 1_A and $x \in \mathcal{M}^B$ is a solution of (1.7); since $N_{A/K}(\varepsilon) \in \mathcal{O}_S^*$ for $\varepsilon \in \mathfrak{U}_{\mathcal{M},B}$, every element of $x\mathfrak{U}_{\mathcal{M},B}$ is a solution of (1.7). If $A = M$ is a finite extension field of K this notion of family of solutions coincides with that for norm form equation (1.3) since then, the K -subalgebras of A containing 1_A are precisely the subfields of M containing K . In [5], Györy proved among other things that the set of solutions of (1.7) is the union of finitely many families.

1.2 Results.

Below, we first recall Gyóry's result on the number of families of solutions of (1.7) and then state our improvement. As before, let K be an algebraic number field, S a finite set of places on K containing all infinite places, $A = M_1 \oplus \cdots \oplus M_t$ where M_1, \dots, M_t are finite extensions of K , and \mathcal{M} a finitely generated (not necessarily free) \mathcal{O}_S -submodule of A . Let $a_i = (\alpha_{i1}, \dots, \alpha_{it})$ ($i = 1, \dots, m$) be a set of generators of \mathcal{M} . Thus,

$$\mathcal{M} = \{x = (l_1(\mathbf{x}), \dots, l_t(\mathbf{x})) : \mathbf{x} \in \mathcal{O}_S^m\}$$

where $l_j(\mathbf{x}) = \alpha_{1j}x_1 + \cdots + \alpha_{mj}x_m$ for $j = 1, \dots, t$, and by (1.6) we have $N_{A/K}(x) = \prod_{j=1}^t N_{M_j/K}(l_j(\mathbf{x}))$. We call d a *denominator* of \mathcal{M} if $d \in K^*$ and if the polynomial $d \prod_{j=1}^t N_{M_j/K}(l_j(\mathbf{X}))$ has its coefficients in \mathcal{O}_S . This notion of denominator is easily shown to be independent of the choice of the generators a_1, \dots, a_m .

We consider eq. (1.7), and impose the following conditions on S , A , \mathcal{M} , β and c :

$$(1.12) \quad \begin{cases} S \text{ has cardinality } s, \\ A \text{ has dimension as a } K\text{-vector space } \sum_{i=1}^t [M_i : K] = r \geq 2, \\ \text{the } K\text{-vector space } V := K\mathcal{M} \text{ has dimension } n \geq 2, \\ \beta \in \mathcal{O}_S \setminus \{0\}, c \text{ is a denominator of } \mathcal{M}. \end{cases}$$

For every finite place v on K , let $\text{ord}_v(\cdot)$ denote the discrete valuation corresponding to v with value group \mathbf{Z} ; recall that $|\cdot|_v = C_v^{-\text{ord}_v(\cdot)}$ for some $C_v > 1$. For $\beta \in K^*$, let $\omega_S(\beta)$ denote the number of $v \notin S$ with $\text{ord}_v(\beta) \neq 0$ and put

$$\psi_1(\beta) := \binom{r}{n-1}^{\omega_S(\beta)} \cdot \prod_{v \notin S} \binom{r \cdot \text{ord}_v(\beta) + n}{n}.$$

Further, let D be the degree over \mathbf{Q} of the normal closure of the composite $M_1 \dots M_t$ over \mathbf{Q} ; thus, $[K : \mathbf{Q}] \leq D \leq (r[K : \mathbf{Q}])!$. Gyóry [5] proved that the set of solutions of (1.7) is contained in some finite union of cosets of unit groups

$$(1.13) \quad x_1 \mathcal{O}_{B_1, S}^* \cup \cdots \cup x_w \mathcal{O}_{B_w, S}^* \text{ with } w \leq (4sD)^{2^{37n} s^6} \cdot \psi_1(\beta),$$

where for $i = 1, \dots, w$, B_i is a K -subalgebra of A with $1_A \in B_i$, $x_i \in A^*$ with $x_i B_i \subset V$, and where the set of solutions of (1.7) contained in $x_i \mathcal{O}_{B_i, S}^*$ is the union of at most $[\mathcal{O}_{B_i, S}^* : \mathfrak{U}_{\mathcal{M}, B_i}]$ (\mathcal{M}, B_i) -families of solutions. This implies an upper bound for the number of families of solutions of (1.7) which depends on n, r, β, s and the indices $[\mathcal{O}_{B_i, S}^* : \mathfrak{U}_{\mathcal{M}, B}]$ (cf. [5], Theorem 3), so ultimately on the module \mathcal{M} . We mention that Voutier [18], Chap. V independently obtained a result similar to (1.13) but only for norm form equation (1.3) and with $K = \mathbf{Q}$, $\beta = 1$.

Gyóry's result can be improved as follows. A K -subalgebra B of A is said to be *S-minimal* if $1_A \in B$, and if for each proper K -subalgebra B' of B with $1_A \in B'$,

the quotient group $\mathcal{O}_{B,S}^*/\mathcal{O}_{B',S}^*$ is infinite. A family of solutions of (1.7) is said to be *reducible* if it is the union of finitely many strictly smaller families of solutions, and *irreducible* otherwise. Put

$$(1.14) \quad \begin{aligned} \psi_2(\beta) &:= \binom{r}{n-1}^{\omega_S(\beta)} \cdot \prod_{v \notin S} \binom{\text{ord}_v(\beta) + n - 1}{n - 1}, \\ e(n) &:= \frac{1}{3}n(n+1)(2n+1) - 2. \end{aligned}$$

Theorem 1. *Assume (1.12). The set of solutions of*

$$(1.7) \quad cN_{A/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M}$$

can be expressed as a finite union of irreducible families of solutions. More precisely, the set of solutions of (1.7) is contained in some finite union of cosets

$$(1.15) \quad x_1\mathcal{O}_{B_1,S}^* \cup \cdots \cup x_w\mathcal{O}_{B_w,S}^* \text{ with } w \leq (2^{33}r^2)^{e(n)s} \cdot \psi_2(\beta)$$

such that for $i = 1, \dots, w$, B_i is an S -minimal K -subalgebra of A , $x_i \in A^$ with $x_i B_i \subset V$, and the set of solutions of (1.7) contained in $x_i\mathcal{O}_{B_i,S}^*$ is the union of at most $[\mathcal{O}_{B_i,S}^* : \mathfrak{U}_{\mathcal{M},B_i}]$ (\mathcal{M}, B_i) -families of solutions which are all irreducible.*

Remark 1. The right-hand side of Györy's bound (1.13) depends doubly exponentially on n and in the worst case that $D = (r[K : \mathbf{Q}])!$ triply exponentially on r , whereas our bound (1.15) depends only polynomially on r and exponentially on n^3 . (1.13) can be better than (1.15) in terms of r only if D is very small compared with r , e.g. if $A = \mathbf{Q}^r$ for some large r . It is likely that, in (1.15), 2^{33} can be improved upon, and that $e(n)$ can be replaced by a linear expression of n .

For some very special type of norm form equation, Voutier succeeded in deriving an upper bound for the number of families of solutions independent of the module \mathcal{M} (see the remark after Corollary 1). It is an open problem whether an explicit bound independent of \mathcal{M} exists in full generality, for equations (1.3) or (1.7).

Remark 2. We can express the set of solutions of (1.7) as a minimal finite union of irreducible families, that is, as a union $\mathcal{F}_1 \cup \cdots \cup \mathcal{F}_g$ where $\mathcal{F}_1, \dots, \mathcal{F}_g$ are irreducible families of solutions, none of which is contained in the union of the others. We claim that every other irreducible family of solutions of (1.7) is contained in one of $\mathcal{F}_1, \dots, \mathcal{F}_g$. In other words $\mathcal{F}_1, \dots, \mathcal{F}_g$ are the maximal irreducible families of solutions of (1.7). Hence Theorem 1 above gives automatically an upper bound for the number of maximal irreducible families. To prove our claim, let \mathcal{G} be an arbitrary irreducible family of solutions of (1.7). Then \mathcal{G} is the union of the sets $\mathcal{G} \cap \mathcal{F}_i$ for $i = 1, \dots, g$ and by Lemma 3 in Section 2, each of these sets is a union of finitely many families. Then one of these families, contained in \mathcal{F}_1 , say, is equal to \mathcal{G} . Hence $\mathcal{G} \subseteq \mathcal{F}_1$.

Remark 3. There is only one way to express the set of solutions of (1.7) as a minimal union of irreducible families, since the families appearing in such a union are the maximal irreducible families of solutions of (1.7).

We also investigate the problem to give an upper bound for the number of K -subalgebras B of A for which (1.7) has (\mathcal{M}, B) -families of solutions. Let again $V = K\mathcal{M}$. Suppose again that $\dim_K A = r$ and $\dim_K V = n$. If x is a solution in \mathcal{M}^B , then $x \in V^B \cap A^*$, where A^* is the unit group of A . Hence (1.7) can have (\mathcal{M}, B) -families of solutions only for those K -subalgebras B of A for which

$$(1.16) \quad 1_A \in B, \quad V^B \cap A^* \neq \emptyset.$$

In [5], Gyóry proved that the number of algebras B with (1.16) is at most n^r . We can improve this as follows:

Theorem 2. *The number of K -subalgebras B of A with (1.16) is at most*

$$(n \max(r - n, 2))^n.$$

We do not know whether the dependence on r is necessary.

We derive some corollaries from Theorem 1. First we specialise Theorem 1 to norm form equation (1.3). Let K, S be as above so that in particular S has cardinality s . Further, let M be a finite extension of K of degree $r \geq 2$, \mathcal{M} a finitely generated \mathcal{O}_S -submodule of M such that the K -vector space $K\mathcal{M}$ has dimension $n \geq 2$, and c, β constants such that $\beta \in \mathcal{O}_S \setminus \{0\}$ and c is a denominator of \mathcal{M} . Then, by applying Theorem 1 with $A = M$, we get at once the following result which improves upon the corresponding results in [5] and [18]:

Corollary 1. *The set of solutions of*

$$(1.3) \quad cN_{M/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M}$$

can be expressed as a finite union of irreducible families of solutions. More precisely, the set of solutions of (1.3) is contained in some finite union of cosets

$$x_1\mathcal{O}_{J_1, S}^* \cup \cdots \cup x_w\mathcal{O}_{J_w, S}^* \text{ with } w \leq (2^{33}r^2)^{e(n)s} \cdot \psi_2(\beta)$$

such that for $i = 1, \dots, w$, J_i is a subfield of M containing K , $x_i \in M^$ is such that $x_i J_i \subset V$, and the set of solutions of (1.3) in $x_i\mathcal{O}_{J_i, S}^*$ is the union of at most $[\mathcal{O}_{J_i, S}^* : \mathfrak{A}_{\mathcal{M}, J_i}]$ (\mathcal{M}, J_i) -families of solutions which are all irreducible.*

As mentioned before, for a very special type of norm form equation Voutier ([18], Theorem V.3) obtained an upper bound for the number of families independent of \mathcal{M} : namely, he proved that if \mathcal{M} is a \mathbf{Z} -module of rank 3 contained in the ring of integers of an algebraic number field M of degree $r > \text{rank } \mathcal{M} = 3$, then the set of solutions of the equation

$$N_{M/\mathbf{Q}}(x) = 1 \text{ in } x \in \mathcal{M}$$

is the union of at most $r^{2^{86}r^2}$ families.

We return to eq. (1.7). In what follows, we consider K as a K -subalgebra of A by indentifying $\alpha \in K$ with $\alpha \cdot 1_A$. The set of solutions of (1.7) can be divided into \mathcal{O}_S^* -cosets $x\mathcal{O}_S^*$. Györy [5], Corollary 2, gave an explicit upper bound for the number of \mathcal{O}_S^* -cosets of solutions of (1.7) in case that this number is finite. We can improve this as follows:

Corollary 2. *Assume (1.12). Suppose that (1.7) has only finitely many \mathcal{O}_S^* -cosets of solutions. Then this number is at most*

$$(2^{33}r^2)^{e(n)s} \cdot \psi_2(\beta) .$$

For $\beta = 1$, this gives the Corollary to Theorem 1 of [4].

Proof. Let B be one of the S -minimal K -subalgebras of A occurring in (1.15). We may assume that (1.7) has an (\mathcal{M}, B) -family of solutions, $x\mathfrak{U}_{\mathcal{M}, B}$, say. By identifying $\varepsilon \in \mathcal{O}_S^*$ with $\varepsilon \cdot 1_A$, we may view \mathcal{O}_S^* as a subgroup of $\mathfrak{U}_{\mathcal{M}, B}$. Let $w \leq \infty$ be the index of \mathcal{O}_S^* in $\mathfrak{U}_{\mathcal{M}, B}$. Then $x\mathfrak{U}_{\mathcal{M}, B}$ is the union of precisely w \mathcal{O}_S^* -cosets. So our assumption implies that w is finite. Therefore, $[\mathcal{O}_{B,S}^* : \mathcal{O}_S^*]$ is finite. Now since B is S -minimal, it follows that $B = K$. So each algebra B_i occurring in (1.15) is equal to K , i.e. $\mathcal{O}_{B_i,S}^* = \mathcal{O}_S^*$, and Corollary 2 follows. \square

In general, it is as yet not effectively decidable whether (1.7) has only finitely many \mathcal{O}_S^* -cosets of solutions. Schmidt [17] Theorem 3, derived an explicit upper bound for the number of solutions of norm form equations over \mathbf{Z} satisfying an effectively decidable non-degeneracy condition. It is possible to give a similar effective non-degeneracy condition for eq. (1.7) as well, which implies that for every $\beta \in \mathcal{O}_S \setminus \{0\}$, the number of \mathcal{O}_S^* -cosets of solutions is finite. Moreover, under that condition we can derive an upper bound for the number of \mathcal{O}_S^* -cosets of solutions with a better dependence on β in that unlike the bound in Corollary 2, it does not depend on the quantities $ord_v(\beta)$ ($v \in M_K \setminus S$) appearing in $\psi_2(\beta)$.

The vector space $V = K\mathcal{M}$ is said to be *non-degenerate* if $V^B \cap A^* = \emptyset$ for every K -subalgebra B of A with $1_A \in B$, $B \neq K$, where A^* is the unit group of A . (1.16) implies that in that case, each algebra B_i occurring in (1.15) is equal to K . Hence the set of solutions of (1.7) is the union of finitely many \mathcal{O}_S^* -cosets.

Corollary 3. *Assume (1.12) and in addition that $V = K\mathcal{M}$ is non-degenerate. Then the set of solutions of (1.7) is the union of at most*

$$(2^{33}r^2)^{e(n)(s+\omega_S(\beta))}$$

\mathcal{O}_S^* -cosets.

Proof. We apply Theorem 1 with $S' := S \cup \{v \notin S : ord_v(\beta) > 0\}$ replacing S . Thus, $\beta \in \mathcal{O}_{S'}^*$. We have to replace s by the cardinality of S' which is $s' := s + \omega_S(\beta)$.

Moreover, in the definition of $\psi_2(\beta)$, S has to be replaced by S' which means that $\psi_2(\beta)$ has to be replaced by 1. Let \mathcal{M}' be the $\mathcal{O}_{S'}$ -module generated by \mathcal{M} . Thus, every solution of (1.7) satisfies

$$(1.7') \quad cN_{A/K}(x) \in \mathcal{O}_{S'}^*, \text{ in } x \in \mathcal{M}' .$$

Clearly, c is a denominator of \mathcal{M}' . Moreover, since V is non-degenerate, the set of solutions of (1.7') is the union of finitely many $\mathcal{O}_{S'}^*$ -cosets. So by Corollary 2, the set of solutions of (1.7'), and hence also the set of solutions of (1.7), is contained in the union of at most $(2^{33}r^2)^{e(n)s'}$ $\mathcal{O}_{S'}^*$ -cosets. Now if any two solutions x_1, x_2 of (1.7) belong to the same $\mathcal{O}_{S'}^*$ -coset then they belong to the same \mathcal{O}_S^* -coset: for if $x_2 = \varepsilon x_1$ with $\varepsilon \in \mathcal{O}_{S'}$, then $\varepsilon^r = cN_{A/K}(x_2)/cN_{A/K}(x_1) \in \mathcal{O}_S^*$, hence $\varepsilon \in \mathcal{O}_S^*$. This proves Corollary 3. \square

2. AN ASYMPTOTIC FORMULA

In this section, we state and prove an asymptotic density result for the collection of \mathcal{O}_S^* -cosets of solutions of equation (1.7), in case that the number of these is infinite. This asymptotic density result is a consequence of (the qualitative part of) Theorem 1.

We recall the definition of absolute (multiplicative) Weil height. Let $\overline{\mathbf{Q}}$ denote the algebraic closure of \mathbf{Q} . Let $\mathbf{x} = (x_1, \dots, x_n) \in \overline{\mathbf{Q}}^n \setminus \{\mathbf{0}\}$. Take any algebraic number field L containing x_1, \dots, x_n , and let $\sigma_1, \dots, \sigma_d$ be the isomorphic embeddings of L into $\overline{\mathbf{Q}}$, where $d = [L : \mathbf{Q}]$. Further, let (x_1, \dots, x_n) denote the fractional ideal with respect to the ring of integers of L generated by x_1, \dots, x_n , and denote by $N_{L/\mathbf{Q}}((x_1, \dots, x_n))$ its norm. Then the absolute Weil height of \mathbf{x} is defined by

$$H(\mathbf{x}) = H(x_1, \dots, x_n) := \left\{ \frac{\prod_{i=1}^d \max(|\sigma_i(x_1)|, \dots, |\sigma_i(x_n)|)}{N_{L/\mathbf{Q}}((x_1, \dots, x_n))} \right\}^{1/d} .$$

It is clear that $H(\mathbf{x})$ does not depend on the choice of L . Further,

$$(2.1) \quad H(\lambda \mathbf{x}) = H(\mathbf{x}) \text{ for } \mathbf{x} \in \overline{\mathbf{Q}}^n \setminus \{\mathbf{0}\}, \lambda \in \overline{\mathbf{Q}}^* .$$

Now let K be an algebraic number field and $A = M_1 \oplus \dots \oplus M_t$, where M_1, \dots, M_t are finite extension fields of K . We define the height $\overline{H}(x)$ of $x = (\xi_1, \dots, \xi_t) \in A$ to be the absolute Weil height of the vector with coordinates consisting of ξ_1, \dots, ξ_t and their conjugates over K , that is, if $\tau_{i,1}, \dots, \tau_{i,r_i}$ with $r_i = [M_i : K]$ are the K -isomorphic embeddings of M_i into $\overline{\mathbf{Q}}$ then we put

$$\overline{H}(x) := H(\tau_{1,1}(\xi_1), \dots, \tau_{1,r_1}(\xi_1), \dots, \tau_{t,1}(\xi_t), \dots, \tau_{t,r_t}(\xi_t)).$$

Note that by (2.1) we have

$$(2.2) \quad \overline{H}(x) = \overline{H}(\lambda x) \text{ for } x \in A \setminus \{0\}, \lambda \in K^* ,$$

i.e. \overline{H} may be viewed as a height on the collection $(A \setminus \{0\})/K^*$ of K^* -cosets xK^* ($x \in A \setminus \{0\}$). This height satisfies

$$(2.3) \quad \#\{x \in (A \setminus \{0\})/K^* : \overline{H}(x) \leq X\} < \infty \text{ for } X > 0 .$$

Namely, by Northcott's theorem [10], [11] we have that for every $d > 0$, $X > 0$, there are, up to multiplication by elements from $\overline{\mathbf{Q}}^*$, only finitely many $\mathbf{x} = (\xi_1, \dots, \xi_n) \in \overline{\mathbf{Q}}^n \setminus \{0\}$ with $H(\mathbf{x}) \leq X$ and $[\mathbf{Q}(\xi_i) : \mathbf{Q}] \leq d$ for $i = 1, \dots, n$. This implies that the set of non-zero elements x of A with $\overline{H}(x) \leq X$ can be divided into finitely many classes, where $x = (\xi_1, \dots, \xi_t)$, $y = (\eta_1, \dots, \eta_t) \in A$ are said to belong to the same class if $(\tau_{1,1}(\xi_1), \dots, \tau_{t,r_t}(\xi_t)) = \alpha(\tau_{1,1}(\eta_1), \dots, \tau_{t,r_t}(\eta_t))$ for some $\alpha \in \overline{\mathbf{Q}}^*$. But clearly, if for instance $\xi_1 \neq 0$, then $\alpha = \tau_{1,1}(\eta_1/\xi_1) = \dots = \tau_{1,r_1}(\eta_1/\xi_1)$ which implies that $\alpha \in K$. So if $x, y \in A \setminus \{0\}$ belong to the same class then they belong to the same K^* -coset.

For a finitely generated abelian group Λ , denote by Λ_{tors} the torsion subgroup of Λ and by $\text{rank } \Lambda$ the rank of the free abelian group $\Lambda/\Lambda_{\text{tors}}$. Let as usual S be a finite set of places on K which contains all infinite places. For a K -subalgebra B of A containing the unit element 1_A of A we put

$$\rho_{B,S} := \text{rank } \mathcal{O}_{B,S}^* / \mathcal{O}_S^* ,$$

where we view \mathcal{O}_S^* as a subgroup of $\mathcal{O}_{B,S}^*$ by identifying $\varepsilon \in \mathcal{O}_S^*$ with $\varepsilon \cdot 1_A$. By a straightforward generalisation of Dirichlet's unit theorem, $\mathcal{O}_{B,S}^*$ is finitely generated, hence $\rho_{B,S}$ is finite.

Let again $\beta, c \in K^*$, and let \mathcal{M} be a finitely generated \mathcal{O}_S -submodule of A such that condition (1.12) holds. For every $X > 0$ we consider the set of solutions of

$$(2.4) \quad cN_{A/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M} \text{ with } \overline{H}(x) \leq X .$$

From (2.2) and $\mathcal{O}_S^* \subset K^*$ it follows that the set of solutions of (2.4) can be divided into \mathcal{O}_S^* -cosets $x\mathcal{O}_S^*$. Denote by $N(X)$ the maximal number of distinct \mathcal{O}_S^* -cosets contained in the set of solutions of (2.4). From (2.3) it follows that $N(X)$ is finite: namely if x, y are solutions of (2.4) with $y = \varepsilon x$ for some $\varepsilon \in K^*$, then $\varepsilon^r = N_{A/K}(y)/N_{A/K}(x) \in \mathcal{O}_S^*$, so x, y belong to the same \mathcal{O}_S^* -coset. In case of norm form equations over \mathbf{Q} , asymptotic formulas for $N(X)$ were derived by Györy and Pethő [6] (in the archimedean case) and Pethő [12] (for an arbitrary finite set of places S); Györy and Pethő [7] and Everest [2] obtained more precise results in certain special cases. From (the qualitative part of) Theorem 1 we derive the following generalisation of Pethő's result [12]:

Corollary 4. *We have*

$$N(X) = \gamma \cdot (\log X)^\rho + O((\log X)^{\rho-1}) \text{ as } X \rightarrow \infty ,$$

where γ is a positive number independent of X and where ρ is the maximum of the numbers $\rho_{B,S}$, taken over all K -subalgebras B of A with $1_A \in B$ for which the equation $cN_{A/K}(x) \in \beta\mathcal{O}_S^*$ in $x \in \mathcal{M}$ has (\mathcal{M}, B) -families of solutions.

We mention that in the case $\mathcal{O}_S = \mathbf{Z}$, Everest and Györy [3] recently obtained some refinements for equations of the form (1.4).

Remark 4. γ , ρ and the constant in the error term are all ineffective. By (1.16), we can estimate ρ from above by the effectively computable number ρ_0 , which is the maximum of the numbers $\rho_{B,S}$, taken over all K -subalgebras B of A with $1_A \in B$, $V^B \cap A^* \neq \emptyset$. Further, using the explicit bound in Theorem 1, one can effectively compute an upper bound for γ ; we shall not work this out.

To derive Corollary 4 we need some lemmas. The first lemma is undoubtedly well-known but we could not find a proof of it in the literature.

Lemma 1. *Let Λ be a finitely generated additive abelian group of rank ρ , and let f be a function from Λ to \mathbf{R} with the following properties:*

$$(2.5) \quad f(x) \geq 0 \text{ for } x \in \Lambda;$$

$$(2.6) \quad f(x+y) \leq f(x) + f(y) \text{ for } x, y \in \Lambda;$$

$$(2.7) \quad f(\lambda x) = \lambda f(x) \text{ for } x \in \Lambda, \lambda \in \mathbf{Z}_{\geq 0};$$

$$(2.8) \quad \text{for every } Y > 0, \text{ the set } \{x \in \Lambda : f(x) \leq Y\} \text{ is finite.}$$

Then

$$(2.9) \quad \#\{x \in \Lambda : f(x) \leq Y\} = \gamma \cdot Y^\rho + O(Y^{\rho-1}) \text{ as } Y \rightarrow \infty$$

where $\gamma = \gamma(\Lambda, f)$ is a positive constant.

Proof. We first assume that $\Lambda = \mathbf{Z}^\rho$. For $x = (\xi_1, \dots, \xi_\rho) \in \mathbf{R}^\rho$ we define the maximum norm $\|x\| := \max(|\xi_1|, \dots, |\xi_\rho|)$. Letting $e_i = (0, \dots, 1, \dots, 0)$ ($i = 1, \dots, \rho$) denote the vector in \mathbf{Z}^ρ with a single 1 on the i -th place, we infer from (2.5)-(2.7) that for $x = (\xi_1, \dots, \xi_\rho)$, $y = (\eta_1, \dots, \eta_\rho) \in \mathbf{Z}^\rho$ we have

$$|f(x) - f(y)| \leq \max(f(x-y), f(y-x)) \leq \sum_{i=1}^{\rho} |\xi_i - \eta_i| \max(f(e_i), f(-e_i)),$$

whence

$$(2.10) \quad |f(x) - f(y)| \leq C \cdot \|x - y\|,$$

where $C := \sum_{i=1}^{\rho} \max(f(e_i), f(-e_i))$.

We extend f to a function on \mathbf{Q}^ρ by putting $f(x) := \lambda^{-1}f(\lambda x)$ for $x \in \mathbf{Q}^\rho$ where λ is the smallest positive integer such that $\lambda x \in \mathbf{Z}^\rho$. This extended f satisfies again (2.5)-(2.7) and (2.10), but now for all $x, y \in \mathbf{Q}^\rho$ and $\lambda \in \mathbf{Q}_{\geq 0}$. Using (2.10) and

taking limits we can extend f to a continuous function $f : \mathbf{R}^\rho \mapsto \mathbf{R}$ which satisfies (2.5)-(2.7) and (2.10) for all $x, y \in \mathbf{R}^\rho$ and $\lambda \in \mathbf{R}_{\geq 0}$.

For $Y > 0$ we define the set $\mathcal{C}_Y := \{x \in \mathbf{R}^\rho : f(x) \leq Y\}$. Since f is continuous, this set is Lebesgue measurable. By (2.7) we have $\mathcal{C}_Y = \{Yx : x \in \mathcal{C}_1\}$. Hence \mathcal{C}_Y has Lebesgue measure $\gamma \cdot Y^\rho$, where γ is the Lebesgue measure of \mathcal{C}_1 . We can cover \mathbf{R}^ρ by the unit cubes $U_z := \{x \in \mathbf{R}^\rho : \|x - z\| \leq \frac{1}{2}\}$ ($z \in \mathbf{Z}^\rho$). These cubes have Lebesgue measure 1, and any two different cubes have at most part of their boundary in common. (2.7) and (2.10) imply that

$$\mathcal{C}_{Y - \frac{1}{2}C} \subseteq \bigcup_{\substack{z \in \mathbf{Z}^\rho \\ f(z) \leq Y}} U_z \subseteq \mathcal{C}_{Y + \frac{1}{2}C} \text{ for } Y \geq \frac{1}{2}C .$$

Now let $n(Y)$ be the number of $z \in \mathbf{Z}^\rho$ with $f(z) \leq Y$. By comparing Lebesgue measures, we get

$$(2.11) \quad \gamma \cdot (Y - \frac{1}{2}C)^\rho \leq n(Y) \leq \gamma \cdot (Y + \frac{1}{2}C)^\rho \text{ for } Y \geq \frac{1}{2}C .$$

From (2.8) it follows that $n(Y)$ is finite; hence γ is finite. Moreover, for Y sufficiently large, $n(Y) > 0$, hence $\gamma > 0$. Now (2.9) follows at once from (2.11). This settles the case $\Lambda = \mathbf{Z}^\rho$.

Now let Λ be an arbitrary additive abelian group. There are $u_1, \dots, u_\rho \in \Lambda$ such that every $x \in \Lambda$ can be expressed uniquely as

$$x = t + \zeta_1 u_1 + \dots + \zeta_\rho u_\rho \text{ with } t \in \Lambda_{\text{tors}}, z = (\zeta_1, \dots, \zeta_\rho) \in \mathbf{Z}^\rho .$$

Put $f'(z) := f(\zeta_1 u_1 + \dots + \zeta_\rho u_\rho)$. (2.6) implies that $f'(z) - f(-t) \leq f(x) \leq f'(z) + f(t)$. Further, (2.7) with $\lambda = 0$ implies that $f(0) = 0$. More generally, (2.7) implies that $f(t) = 0$ for $t \in \Lambda_{\text{tors}}$ since for such t there is a positive integer λ with $\lambda t = 0$. Hence $f(x) = f'(z)$ for $x \in \Lambda$. Clearly, f' and \mathbf{Z}^ρ satisfies (2.5)-(2.8). So by what we proved above we have

$$\#\{z \in \mathbf{Z}^\rho : f'(z) \leq Y\} = \gamma' Y^\rho + O(Y^{\rho-1}) \text{ as } Y \rightarrow \infty$$

with some positive γ' . From this, one deduces easily that (2.9) holds with $\gamma = \gamma' \cdot \#\Lambda_{\text{tors}}$. This completes the proof of Lemma 1. \square

For a subset \mathcal{F} of A with the property that for each $x \in \mathcal{F}$ the coset $x\mathcal{O}_S^*$ is contained in \mathcal{F} , we denote by $N_{\mathcal{F}}(X)$ the maximal number of distinct \mathcal{O}_S^* -cosets $x\mathcal{O}_S^*$ with $x \in \mathcal{F}$ and $\overline{H}(x) \leq X$.

Lemma 2. *Let $\mathcal{F} = x\mathcal{M}_{\mathcal{M}, B}$ be a family of solutions of (1.7), where B is a K -subalgebra of A containing 1_A and $x \in \mathcal{M}^B$. Then for some positive real γ depending only on \mathcal{M} and B we have*

$$(2.12) \quad N_{\mathcal{F}}(X) = \gamma(\log X)^{\rho_{B,S}} + O((\log X)^{\rho_{B,S}-1}) \text{ as } X \rightarrow \infty .$$

Proof. We use the following properties of the absolute Weil height which are straightforward consequences of its definition:

$$(2.13) \quad \begin{cases} H(\mathbf{x}) \geq 1 \text{ for } \mathbf{x} \in \overline{\mathbf{Q}}^n \setminus \{\mathbf{0}\}, \\ H(x_1 y_1, \dots, x_n y_n) \leq H(x_1, \dots, x_n) H(y_1, \dots, y_n) \\ \text{for } x_1, \dots, x_n, y_1, \dots, y_n \in \overline{\mathbf{Q}}, \\ H(x_1^\lambda, \dots, x_n^\lambda) = H(x_1, \dots, x_n)^\lambda \text{ for } x_1, \dots, x_n \in \overline{\mathbf{Q}}, \lambda \in \mathbf{Z}_{\geq 0}. \end{cases}$$

Let $\mathfrak{U} := \mathfrak{U}_{\mathcal{M}, B}$ and $\rho_0 := \rho_{B, S}$. Since \mathfrak{U} has finite index in $\mathcal{O}_{B, S}^*$, the factor group $\mathfrak{U}/\mathcal{O}_S^*$ has rank ρ_0 . We apply Lemma 1 to $\Lambda = \mathfrak{U}/\mathcal{O}_S^*$ and $f = \log \overline{H}$. By (2.2), f is well-defined on Λ . Further, (2.13) implies (2.5)-(2.7), and (2.8) follows from (2.3) and the fact that $\mathfrak{U}/\mathcal{O}_S^* = \mathfrak{U}/(K^* \cap \mathfrak{U})$ may be viewed as a subgroup of A^*/K^* . It follows that

$$(2.14) \quad N_{\mathfrak{U}}(X) = \gamma(\log X)^{\rho_0} + O((\log X)^{\rho_0-1}) \text{ as } X \rightarrow \infty$$

for some positive constant γ . By (2.13) we have $c_1 \overline{H}(xu) \leq \overline{H}(u) \leq c_2 \overline{H}(xu)$ for $u \in \mathfrak{U}$, where $c_1 = \overline{H}(x)^{-1}$ and $c_2 = \overline{H}(x^{-1})$, and this implies that $N_{\mathfrak{U}}(c_2^{-1} X) \leq N_{x\mathfrak{U}}(X) \leq N_{\mathfrak{U}}(c_1^{-1} X)$. Now Lemma 2 follows from (2.14) and the fact that both $(\log(c_1^{-1} X))^{\rho_0}$ and $(\log(c_2^{-1} X))^{\rho_0}$ differ from $(\log X)^{\rho_0}$ by at most $O((\log X)^{\rho_0-1})$. \square

Lemma 3. *For any two K -subalgebras B_1, B_2 of A containing 1_A , the intersection of an (\mathcal{M}, B_1) -family and an (\mathcal{M}, B_2) -family is the union of at most finitely many $(\mathcal{M}, B_1 \cap B_2)$ -families.*

Proof. Let $\mathcal{G}_i = x_i \mathfrak{U}_{\mathcal{M}, B_i}$ with $x_i \in \mathcal{M}^{B_i}$ for $i = 1, 2$ be the two families of solutions and put $B := B_1 \cap B_2$. Let $x_0 \in \mathcal{G}_1 \cap \mathcal{G}_2$. Then $x_0 \in \mathcal{M}^{B_1} \cap \mathcal{M}^{B_2}$. From definition (1.10) it follows easily that $\mathcal{M}^{B_i} \subseteq \mathcal{M}^B$ for $i = 1, 2$. Therefore, $x_0 \in \mathcal{M}^B$. Further, we have $\mathcal{G}_i = x_0 \mathfrak{U}_{\mathcal{M}, B_i}$ for $i = 1, 2$, hence $\mathcal{G}_1 \cap \mathcal{G}_2 = x_0 (\mathfrak{U}_{\mathcal{M}, B_1} \cap \mathfrak{U}_{\mathcal{M}, B_2})$. We claim that $\mathfrak{U}_{\mathcal{M}, B}$ is a subgroup of finite index in $\mathfrak{U}_{\mathcal{M}, B_1} \cap \mathfrak{U}_{\mathcal{M}, B_2}$; then it follows at once that $\mathcal{G}_1 \cap \mathcal{G}_2$ is the union of finitely many families $y \mathfrak{U}_{\mathcal{M}, B}$ with $y \in \mathcal{M}^B$. To prove the claim, let $\varepsilon \in \mathfrak{U}_{\mathcal{M}, B}$ and take $i \in \{1, 2\}$. Then $\varepsilon \in B \subseteq B_i$, whence by (1.10), $\varepsilon \mathcal{M}^{B_i} \subseteq V^{B_i}$ where $V = K\mathcal{M}$. Further, by (1.11) we have $\varepsilon \mathcal{M}^{B_i} \subseteq \varepsilon \mathcal{M}^B = \mathcal{M}^B \subseteq \mathcal{M}$. Therefore, by (1.10) $\varepsilon \mathcal{M}^{B_i} \subseteq \mathcal{M}^{B_i}$. Similarly, we find $\varepsilon^{-1} \mathcal{M}^{B_i} \subseteq \mathcal{M}^{B_i}$. Hence $\varepsilon \mathcal{M}^{B_i} = \mathcal{M}^{B_i}$, i.e. $\varepsilon \in \mathfrak{U}_{\mathcal{M}, B_i}$ for $i = 1, 2$. So $\mathfrak{U}_{\mathcal{M}, B} \subseteq \mathfrak{U}_{\mathcal{M}, B_1} \cap \mathfrak{U}_{\mathcal{M}, B_2}$. Now our claim follows from the fact that both groups have finite index in $\mathcal{O}_{B, S}^* = \mathcal{O}_{B_1, S}^* \cap \mathcal{O}_{B_2, S}^*$. \square

Proof of Corollary 4. By Theorem 1, the set of solutions of (1.7) can be expressed as

$$(2.15) \quad \mathcal{F}_1 \cup \dots \cup \mathcal{F}_p$$

where for each i , \mathcal{F}_i is an (\mathcal{M}, B_i) -family of solutions of (1.7) for some K -subalgebra B_i of A containing 1_A . For a tuple $I = \{i_1 < \dots < i_t\}$ of integers from $\{1, \dots, p\}$, let $B_I := B_{i_1} \cap \dots \cap B_{i_t}$, $\mathcal{F}_I := \mathcal{F}_{i_1} \cap \dots \cap \mathcal{F}_{i_t}$, and $N_I(X)$ the number of cosets $x\mathcal{O}_S^*$ with $x \in \mathcal{F}_I$ and $\overline{H}(x) \leq X$. Put $\rho_1 := \max\{\rho_{B_i, S} : i = 1, \dots, p\}$. Thus, $\rho_{B_I, S} \leq \rho_1$ for each tuple I as above. Lemma 3 implies that for each I , \mathcal{F}_I is the union of finitely many (\mathcal{M}, B_I) -families. So by Lemma 2 we have

$$N_I(X) = \gamma_I(\log X)^{\rho_1} + O((\log X)^{\rho_1-1}) \text{ as } X \rightarrow \infty$$

where $\gamma_I = 0$ if $\rho_{B_I, S} < \rho_1$. Note that $\gamma_i > 0$ for at least one $i \in \{1, \dots, p\}$. Now by (2.15) and the rule of inclusion and exclusion we have

$$N(X) = \sum_{i=1}^p N_i(X) - \sum_{\#I=2} N_I(X) + \sum_{\#I=3} N_I(X) - \dots,$$

hence

$$N(X) = \gamma(\log X)^{\rho_1} + O((\log X)^{\rho_1-1}) \text{ as } X \rightarrow \infty$$

where

$$\gamma = \sum_{i=1}^p \gamma_i - \sum_{\#I=2} \gamma_I + \sum_{\#I=3} \gamma_I - \dots.$$

Since $N(X) \geq N_i(X)$ for $i = 1, \dots, p$ we have $\gamma \geq \gamma_i$ for $i = 1, \dots, p$, hence $\gamma > 0$. Lemma 2 implies that (1.7) does not have any family of solutions $x\mathcal{U}_{\mathcal{M}, B}$ with $\rho_{B, S} > \rho_1$; therefore, $\rho_1 = \rho$. This completes the proof of Corollary 4. \square

3. REDUCTION TO $\mathcal{O}_{A, S}^*$ -COSETS

Let K be an algebraic number field, and let $S, M_1, \dots, M_t, A = M_1 \oplus \dots \oplus M_t, \mathcal{M}$ be as in Section 1.2. Further, let $s = \#S, r = \dim_K A \geq 2, n = \dim_K K\mathcal{M} \geq 2, c, \beta$ be as in (1.12). For $x \in A$, we define the coset $x\mathcal{O}_{A, S}^* = \{\varepsilon x : \varepsilon \in \mathcal{O}_{A, S}^*\}$. In this section we prove Lemma 4 below which is in fact an improvement of Lemma 5 of [5].

Lemma 4. *The set of solutions of*

$$(1.7) \quad cN_{A/K}(x) \in \beta\mathcal{O}_S^* \text{ in } x \in \mathcal{M}$$

is contained in some union $x_1\mathcal{O}_{A, S}^ \cup \dots \cup x_{t_1}\mathcal{O}_{A, S}^*$ where $t_1 \leq \psi_2(\beta)$ and where for $j = 1, \dots, t_1, x_j \in \mathcal{M}$ is a solution of (1.7).*

We prove this by slightly refining some arguments of Schmidt [17]. In the proof of Lemma 4 we need some further lemmas. We first recall some lemmas from [17]. Let E be a field endowed with a non-archimedean additive valuation V (i.e. $V(xy) = V(x) + V(y), V(x+y) \geq \min(V(x), V(y))$ for $x, y \in E, V(0) = \infty$, and there is an $x \in E$ with $V(x) \neq 0, V(x) \neq \infty$). For $\mathbf{z} = (z_1, \dots, z_n) \in E^n$, put $V(\mathbf{z}) = \min(V(z_1), \dots, V(z_n))$. Further, let L_1, \dots, L_r be $r \geq n$ linear forms in n variables with coefficients in E .

Lemma 5. *Let $\mathbf{z} \in E^n$ with $\mathbf{z} \neq \mathbf{0}$. There is a subset \mathcal{S} of $\{1, \dots, r\}$ of cardinality $n - 1$ such that every $\mathbf{z}' \in E^n$ with*

$$V(\mathbf{z}') \geq V(\mathbf{z}), \quad V(L_i(\mathbf{z}')) \geq V(L_i(\mathbf{z})) \quad \text{for } i \in \mathcal{S}$$

satisfies

$$V(L_i(\mathbf{z}')) \geq V(L_i(\mathbf{z})) \quad \text{for } i = 1, \dots, r.$$

Proof. This is precisely Lemma 13 of [17], except that that Lemma has the additional condition $V(\mathbf{z}) = 0$. Suppose that $V(\mathbf{z}) \neq 0$. Let $\lambda \in E$ be such that $V(\lambda) = V(\mathbf{z})$ and put $\mathbf{z}_1 := \lambda^{-1}\mathbf{z}$. Then $V(\mathbf{z}_1) = 0$. Now Lemma 5 follows at once from Lemma 13 of [17] applied to \mathbf{z}_1 , on observing that $V(L_i(\mathbf{z}_1)) = V(L_i(\mathbf{z})) - V(\lambda)$ for $i = 1, \dots, r$. \square

We call the subset \mathcal{S} related to \mathbf{z} as in Lemma 5 an *anchor* for \mathbf{z} .

Lemma 6. *Let d_1, \dots, d_r be positive rational numbers, γ a real and \mathcal{S} a subset of $\{1, \dots, r\}$ of cardinality $n - 1$. Put*

$$\mathcal{T}(\mathcal{S}) := \left\{ \mathbf{z} \in E^n : \sum_{i=1}^r d_i V(L_i(\mathbf{z})) = \gamma, \mathcal{S} \text{ is an anchor for } \mathbf{z} \right\}.$$

Then for any $\mathbf{z}_1, \mathbf{z}_2 \in \mathcal{T}(\mathcal{S})$ with $V(L_i(\mathbf{z}_1)) = V(L_i(\mathbf{z}_2))$ for $i \in \mathcal{S}$ we have that $V(L_i(\mathbf{z}_1)) = V(L_i(\mathbf{z}_2))$ for $i = 1, \dots, r$.

Proof. Let $\mathbf{z}_1, \mathbf{z}_2 \in \mathcal{T}(\mathcal{S})$ with $V(L_i(\mathbf{z}_1)) = V(L_i(\mathbf{z}_2))$ for $i \in \mathcal{S}$. We may assume without loss of generality that $V(\mathbf{z}_2) \geq V(\mathbf{z}_1)$. Then by Lemma 5 we have $V(L_i(\mathbf{z}_2)) \geq V(L_i(\mathbf{z}_1))$ for $i = 1, \dots, r$. Together with $\sum_{i=1}^r d_i V(L_i(\mathbf{z}_j)) = \gamma$ for $j = 1, 2$ this implies that $V(L_i(\mathbf{z}_2)) = V(L_i(\mathbf{z}_1))$ for $i = 1, \dots, r$. \square

As before, if we express an element of A as a t -tuple (ξ_1, \dots, ξ_t) , say, then it is implicitly assumed that $\xi_i \in M_i$ for $i = 1, \dots, t$. Fix $v \in M_K \setminus S$. For $i = 1, \dots, t$, let $w_{i,1}, \dots, w_{i,g_i}$ denote the places on M_i which lie above v , and denote by e_{ij}, f_{ij} the ramification index and residue class degree, respectively, of w_{ij} over v . Let \overline{K} denote the algebraic closure of K . Choose a continuation of ord_v to \overline{K} and denote this also by ord_v ; then ord_v assumes its values in \mathbf{Q} . For $i = 1, \dots, t$ let \mathcal{E}_i denote the collection of K -isomorphic embeddings of M_i into \overline{K} ; then \mathcal{E}_i can be expressed as a disjoint union,

$$\mathcal{E}_i = \mathcal{E}_{i1} \cup \dots \cup \mathcal{E}_{ig_i} \quad \text{with } \#\mathcal{E}_{ij} = e_{ij}f_{ij} \quad \text{for } j = 1, \dots, g_i$$

such that for $j = 1, \dots, g_i$

$$(3.1) \quad \text{ord}_{w_{ij}}(\alpha) = e_{ij} \text{ord}_v(\sigma(\alpha)) \quad \text{for } \alpha \in M_i, \sigma \in \mathcal{E}_{ij}.$$

Lemma 7. *There are integers c_{ij} ($i = 1, \dots, t, j = 1, \dots, g_i$) and u_v with $u_v \leq \text{ord}_v(\beta)$ such that for every solution $x = (\xi_1, \dots, \xi_t) \in \mathcal{M}$ of (1.7) we have*

$$(3.2) \quad \text{ord}_{w_{ij}}(\xi_i) - c_{ij} \geq 0 \text{ for } i = 1, \dots, t, j = 1, \dots, g_i,$$

$$(3.3) \quad \sum_{i=1}^t \sum_{j=1}^{g_i} f_{ij} \{ \text{ord}_{w_{ij}}(\xi_i) - c_{ij} \} = u_v.$$

Proof. Let $\{\mathbf{a}_k = (\alpha_{k1}, \dots, \alpha_{kt}) : k = 1, \dots, m\}$ be a set of generators of \mathcal{M} as an \mathcal{O}_S -module. Define the integers

$$(3.4) \quad c_{ij} = \min\{\text{ord}_{w_{ij}}(\alpha_{ki}) : k = 1, \dots, m\} \text{ for } i = 1, \dots, t, j = 1, \dots, g_i.$$

Let $x = (\xi_1, \dots, \xi_t) \in \mathcal{M}$ be a solution of (1.7). Then $x = \sum_{k=1}^m \beta_k \mathbf{a}_k$ for certain $\beta_1, \dots, \beta_m \in \mathcal{O}_S$. Since the place w_{ij} lies above $v \in M_K \setminus S$, we have $\text{ord}_{w_{ij}}(\beta_k) \geq 0$ for $i = 1, \dots, t, j = 1, \dots, g_i$. Together with $\xi_i = \sum_{k=1}^m \beta_k \alpha_{ki}$ for $i = 1, \dots, t$ and (3.4), this implies $\text{ord}_{w_{ij}}(\xi_i) \geq c_{ij}$ for $i = 1, \dots, t, j = 1, \dots, g_i$. This proves (3.2).

We now prove (3.3) for some u_v . By assumption, c is a denominator for \mathcal{M} , i.e.

$$c \prod_{i=1}^t N_{M_i/K}(\alpha_{1i}X_1 + \dots + \alpha_{mi}X_m) \in \mathcal{O}_S[X_1, \dots, X_m].$$

Since $x = (\xi_1, \dots, \xi_t)$ is a solution of (1.7) we have $c \prod_{i=1}^t N_{M_i/K}(\xi_i) \in \beta \mathcal{O}_S^*$, so

$$(3.5) \quad F(\mathbf{X}) = \beta \prod_{i=1}^t N_{M_i/K} \left(\sum_{k=1}^m \frac{\alpha_{ki}}{\xi_i} X_k \right) \in \mathcal{O}_S[X_1, \dots, X_m].$$

For a polynomial $P(\mathbf{X}) \in \overline{K}[X_1, \dots, X_m]$ denote by $\text{ord}_v(P)$ the minimum of the numbers $\text{ord}_v(\alpha)$ for all coefficients α of P . By Gauss' lemma (cf. [8], p.55, Prop.2.1) we have $\text{ord}_v(PQ) = \text{ord}_v(P) + \text{ord}_v(Q)$ for $P, Q \in \overline{K}[X_1, \dots, X_m]$. By applying this to (3.5) we obtain

$$\begin{aligned} 0 &\leq \text{ord}_v(F) = \text{ord}_v(\beta) + \sum_{i=1}^t \sum_{\sigma \in \mathcal{E}_i} \min_{1 \leq k \leq m} \text{ord}_v(\sigma(\alpha_{ki}/\xi_i)) \\ &= \text{ord}_v(\beta) + \sum_{i=1}^t \sum_{j=1}^{g_i} \sum_{\sigma \in \mathcal{E}_{ij}} \min_{1 \leq k \leq m} \text{ord}_v(\sigma(\alpha_{ki}/\xi_i)) \\ &= \text{ord}_v(\beta) + \sum_{i=1}^t \sum_{j=1}^{g_i} f_{ij} \min_{1 \leq k \leq m} \text{ord}_{w_{ij}}(\alpha_{ki}/\xi_i) \text{ by (3.1)} \\ &= \text{ord}_v(\beta) + \sum_{i=1}^t \sum_{j=1}^{g_i} f_{ij} \{ c_{ij} - \text{ord}_{w_{ij}}(\xi_i) \} \text{ by (3.4)}. \end{aligned}$$

This implies (3.3) with $u_v = \text{ord}_v(\beta) - \text{ord}_v(F)$. \square

Lemma 8. *If $x = (\xi_1, \dots, \xi_t)$ runs through the set of solutions of (1.7), then the tuple $\psi_v(x) := (\text{ord}_{w_{ij}}(\xi_i) : i = 1, \dots, t, j = 1, \dots, g_i)$ runs through a set of cardinality at most $\binom{r}{n-1} \binom{\text{ord}_v(\beta) + n - 1}{n-1}$.*

Proof. Let

$$\mathcal{O}_v := \{y \in K : \text{ord}_v(y) \geq 0\}, \quad \mathcal{M}_v := \mathcal{M}\mathcal{O}_v$$

be the local ring at v , and the localisation of \mathcal{M} at v , respectively. We note that $\mathcal{O}_S \subset \mathcal{O}_v$ and $\mathcal{M} \subset \mathcal{M}_v$. Since \mathcal{O}_v is a principal ideal domain, the \mathcal{O}_v -module \mathcal{M}_v is free of rank $n = \dim_K K\mathcal{M}$. Let $\{\mathbf{a}_k = (\alpha_{k1}, \dots, \alpha_{kt}) : k = 1, \dots, n\}$ be an \mathcal{O}_v -basis of \mathcal{M}_v . Further, let $x = (\xi_1, \dots, \xi_t) \in \mathcal{M}$ be a solution of (1.7). Then $x = z_1\mathbf{a}_1 + \dots + z_n\mathbf{a}_n$ for some vector $\mathbf{z} = (z_1, \dots, z_n) \in \mathcal{O}_v^n$ which is uniquely determined by x . For each $i \in \{1, \dots, t\}$ and each $\sigma \in \mathcal{E}_i$ (the collection of K -isomorphic embeddings of M_i into \overline{K}) define the linear form $L_{i\sigma}(\mathbf{z}) := \sigma(\alpha_{1i})z_1 + \dots + \sigma(\alpha_{ni})z_n$. Thus

$$(3.6) \quad \sigma(\xi_i) = L_{i\sigma}(\mathbf{z}) \text{ for } i = 1, \dots, t, \sigma \in \mathcal{E}_i.$$

Recall that $\sum_{i=1}^t [M_i : K] = r$. Let L_1, \dots, L_r be the linear forms $L_{i\sigma}$ ($i = 1, \dots, t, \sigma \in \mathcal{E}_i$) in some order. For $i = 1, \dots, t, j = 1, \dots, g_i$, let

$$\mathcal{F}_{ij} = \{k \in \{1, \dots, r\} : L_k = L_{i\sigma} \text{ for some } \sigma \in \mathcal{E}_i\},$$

where the set \mathcal{E}_{ij} is defined by (3.1). Then by (3.1), (3.6),

$$(3.7) \quad \begin{aligned} \text{ord}_{w_{ij}}(\xi_i) &= e_{ij} \text{ord}_v(\sigma(\xi_i)) = e_{ij} \text{ord}_v(L_k(\mathbf{z})) \\ &\text{for } i = 1, \dots, t, j = 1, \dots, g_i, k \in \mathcal{F}_{ij}. \end{aligned}$$

We apply Lemma 6 with $E = \overline{K}$ and $V = \text{ord}_v$. Let $\mathcal{S}_x \subset \{1, \dots, r\}$ be an anchor for \mathbf{z} in the sense of Lemma 5. Then \mathcal{S}_x has cardinality $n - 1$, and the tuple $(\text{ord}_v(L_k(\mathbf{z})) : k = 1, \dots, r)$ is uniquely determined by \mathcal{S}_x and the $(n - 1)$ -tuple $(\text{ord}_v(L_k(\mathbf{z})) : k \in \mathcal{S}_x)$. Let

$$\mathcal{S}'_x = \{(i, j) : 1 \leq i \leq t, 1 \leq j \leq g_i, \mathcal{F}_{ij} \cap \mathcal{S}_x \neq \emptyset\}.$$

Now (3.7) implies that once \mathcal{S}_x is given, the tuple $(\text{ord}_{w_{ij}}(\xi_i) : (i, j) \in \mathcal{S}'_x)$ determines uniquely $(\text{ord}_v(L_k(\mathbf{z})) : k \in \mathcal{S}_x)$, the latter determines uniquely $(\text{ord}_v(L_k(\mathbf{z})) : k = 1, \dots, r)$ and this last tuple determines uniquely $(\text{ord}_{w_{ij}}(\xi_i) : i = 1, \dots, t, j = 1, \dots, g_i) = \psi_v(x)$, again by (3.7). We conclude that $\psi_v(x)$ is determined uniquely by \mathcal{S}_x and the tuple $(\text{ord}_{w_{ij}}(\xi_i) : (i, j) \in \mathcal{S}'_x)$.

By Lemma 7 there are integers c_{ij} ($i = 1, \dots, t, j = 1, \dots, g_i$) such that $\text{ord}_{w_{ij}}(\xi_i) - c_{ij} \geq 0$ for $(i, j) \in \mathcal{S}'_x$ and

$$(3.8) \quad \sum_{(i,j) \in \mathcal{S}'_x} \{\text{ord}_{w_{ij}}(\xi_i) - c_{ij}\} \leq \sum_{i=1}^t \sum_{j=1}^{g_i} f_{ij} \{\text{ord}_{w_{ij}}(\xi_i) - c_{ij}\} \leq \text{ord}_v(\beta).$$

The set \mathcal{S}'_x has cardinality $\leq n-1$, since \mathcal{S}_x has cardinality $n-1$ and the sets \mathcal{F}_{ij} are pairwise disjoint. Given the set \mathcal{S}_x , (3.8) implies that for the tuple $(\text{ord}_{w_{ij}}(\xi_i) : (i, j) \in \mathcal{S}'_x)$ we have at most $\binom{\text{ord}_v(\beta) + \#\mathcal{S}'_x}{\#\mathcal{S}'_x} \leq \binom{\text{ord}_v(\beta) + n - 1}{n-1}$ possibilities. Moreover, as \mathcal{S}_x is a subset of $\{1, \dots, r\}$ of cardinality $n-1$, we have at most $\binom{r}{n-1}$ possibilities for \mathcal{S}_x . This proves Lemma 8. \square

Proof of Lemma 4. For $x = (\xi_1, \dots, \xi_t) \in A$ define the tuple of integers $\psi(x) := (\text{ord}_{w_i}(\xi_i) : i = 1, \dots, t, w_i \nmid S)$ where ' $w_i \nmid S$ ' indicates that w_i runs through all places on M_i not lying above a place in S . ψ is an additive homomorphism on A^* with kernel $\mathcal{O}_{A,S}^*$, since $x = (\xi_1, \dots, \xi_t) \in \mathcal{O}_{A,S}^* \iff \xi_i \in \mathcal{O}_{M_i,S}^*$ for $i = 1, \dots, t \iff \text{ord}_{w_i}(\xi_i) = 0$ for $i = 1, \dots, t, w_i \nmid S$. In particular, for $x_1, x_2 \in A^*$ we have $\psi(x_1) = \psi(x_2) \iff x_1 \mathcal{O}_{A,S}^* = x_2 \mathcal{O}_{A,S}^*$.

Now $\psi(x)$ can be obtained by combining all tuples $\psi_v(x)$ ($v \in M_K \setminus S$) from Lemma 8. Hence if x runs through all solutions of (1.7), then $\psi(x)$ runs through a set of cardinality at most

$$\prod_{v \in M_K \setminus S} \binom{r}{n-1} \binom{\text{ord}_v(\beta) + n - 1}{n-1} = \psi_2(\beta).$$

This completes the proof of Lemma 4. \square

4. PROOF OF THEOREM 1

Let $K, S, s = \#S, M_1, \dots, M_t, A = M_1 \oplus \dots \oplus M_t, r = \dim_K A \geq 2, \mathcal{M}, n = \dim_K K\mathcal{M}, c, \beta$ be as in (1.12). Further, put $V := K\mathcal{M}$. By Lemma 4, the set of solutions of (1.7) is contained in some finite union of $\mathcal{O}_{A,S}^*$ -cosets. For the moment, we consider only the solutions of (1.7) in a fixed $\mathcal{O}_{A,S}^*$ -coset $x_0 \mathcal{O}_{A,S}^*$. More generally, we deal with elements of the set

$$(4.1) \quad V \cap x_0 \mathcal{O}_{A,S}^*$$

where x_0 is a fixed element of A^* . As before, we view K as a K -subalgebra of A by identifying $\alpha \in K$ with $\alpha 1_A = (\alpha, \dots, \alpha)$ (r times).

Lemma 9. *Let $B = \{a \in A : aV \subseteq V\}$ be the algebra of scalars of V . Suppose that $n \geq 2$ and that the quotient group $\mathcal{O}_{B,S}^* / \mathcal{O}_S^*$ is finite. Then there are proper K -linear subspaces Y_1, \dots, Y_{t_2} of V such that*

$$V \cap x_0 \mathcal{O}_{A,S}^* \subseteq Y_1 \cup \dots \cup Y_{t_2} \text{ with } t_2 \leq (2^{66} r^4)^{n^2 s}.$$

Proof. We assume that $x_0 = 1$; this is no loss of generality since if $x_0 \neq 1$, we may prove Lemma 9 with $x_0^{-1} V \cap \mathcal{O}_{A,S}^*$ replacing $V \cap x_0 \mathcal{O}_{A,S}^*$. We want to apply Lemma 16 of [4] and for this purpose we must introduce some notation.

For $i = 1, \dots, t$, let $\tau_{i,1}, \dots, \tau_{i,r_i}$ ($r_i = [M_i : K]$) be the K -isomorphic embeddings of M_i into \overline{K} and define the map $f : A \mapsto \overline{K}^r$ by

$$f(x) := (\tau_{1,1}(\xi_1), \dots, \tau_{1,r_1}(\xi_1), \dots, \tau_{t,1}(\xi_t), \dots, \tau_{t,r_t}(\xi_t)) \text{ for } x = (\xi_1, \dots, \xi_t) \in A.$$

Thus, $f(x) = (x_1, \dots, x_r) \in \overline{K}^r$. Let G denote the Galois group of \overline{K}/K . Clearly, for $\sigma \in G$, $i = 1, \dots, t$, $\sigma \circ \tau_{i,1}, \dots, \sigma \circ \tau_{i,r_i}$ is a permutation of $\tau_{i,1}, \dots, \tau_{i,r_i}$. This implies that there is an action by G on $\{1, \dots, r\}$ attaching to each $\sigma \in G$ a permutation $(\sigma(1), \dots, \sigma(r))$ of $(1, \dots, r)$ such that for $x \in A$ we have

$$\sigma(x_i) = x_{\sigma(i)} \text{ for } i = 1, \dots, r, \sigma \in G,$$

where $(x_1, \dots, x_r) = f(x)$. Define the K -algebra

$$\Lambda = \{\mathbf{x} = (x_1, \dots, x_r) \in \overline{K}^r : \sigma(x_i) = x_{\sigma(i)} \text{ for } i = 1, \dots, r, \sigma \in G\}.$$

Then f is an injective K -homomorphism from A to Λ . For instance from Lemma 2 of [4] it follows that K -linearly independent vectors of Λ are also \overline{K} -linearly independent; so $\dim_K \Lambda \leq r = \dim_K A$. It follows that f is also surjective, i.e. a K -algebra isomorphism from A to Λ . Let $\overline{\mathcal{O}}_S$ denote the integral closure of \mathcal{O}_S in \overline{K} , $\overline{\mathcal{O}}_S^*$ the unit group of $\overline{\mathcal{O}}_S$, and $\overline{\mathcal{O}}_S^{*r}$ the r -fold cartesian product of this unit group. It is easy to verify that

$$(4.2) \quad f(\mathcal{O}_{A,S}^*) = \Lambda \cap (\overline{\mathcal{O}}_S^{*r}).$$

A symmetric partition of $\{1, \dots, r\}$ is a collection of sets $\mathcal{P} = \{P_1, \dots, P_q\}$ such that $P_1 \cup \dots \cup P_q = \{1, \dots, r\}$, $P_i \cap P_j = \emptyset$ for $1 \leq i < j \leq q$ and such that for each $P \in \mathcal{P}$, $\sigma \in G$, the set $\sigma(P) = \{\sigma(k) : k \in P\}$ belongs also to \mathcal{P} . To a symmetric partition \mathcal{P} we attach the K -subalgebra of Λ ,

$$\Lambda_{\mathcal{P}} = \{\mathbf{x} = (x_1, \dots, x_r) \in \Lambda : x_i = x_j \text{ for each pair of indices } i, j \text{ belonging to the same set of } \mathcal{P}\}.$$

Let $W := f(V)$ and let \mathcal{P} be a symmetric partition of $\{1, \dots, r\}$ such that

$$(4.3) \quad xW \subseteq W \text{ for } x \in \Lambda_{\mathcal{P}}.$$

Let $\tilde{B} := f^{-1}(\Lambda_{\mathcal{P}})$. Then \tilde{B} is a K -subalgebra of B . Hence $\mathcal{O}_{\tilde{B},S}^*/\mathcal{O}_S^*$ (with $\varepsilon \in \mathcal{O}_S^*$ identified with $(\varepsilon, \dots, \varepsilon)$ (t times)) is finite. Now (4.2) implies that f maps $\mathcal{O}_{\tilde{B},S}^*$ to $\mathcal{O}_{\mathcal{P},S}^* := \Lambda_{\mathcal{P}} \cap (\overline{\mathcal{O}}_S^*)^r$. Further, f maps \mathcal{O}_S^* to $f(\mathcal{O}_S^*) := \{(\varepsilon, \dots, \varepsilon) \text{ (} r \text{ times)} : \varepsilon \in \mathcal{O}_S^*\}$. Hence

$$(4.4) \quad \mathcal{O}_{\mathcal{P},S}^*/f(\mathcal{O}_S^*) \text{ is finite.}$$

Now let \mathcal{P} be the symmetric partition specified in the statement of Lemma 16 of [4]. This \mathcal{P} satisfies (4.3), hence (4.4) and so the condition of Lemma 16 of [4] is

satisfied. Therefore, according to Lemma 16 of [4], the set $W \cap (\overline{\mathcal{O}}_S^*)^r$ is contained in some union $W_1 \cup \dots \cup W_{t_2}$ of proper linear subspaces of W with $t_2 \leq (2^{66}r^4)^{n^2s}$. By (4.2) we have $V \cap \mathcal{O}_{A,S}^* = f^{-1}(W \cap (\overline{\mathcal{O}}_S^*)^r)$. Hence $V \cap \mathcal{O}_{A,S}^* \subseteq Y_1 \cap \dots \cap Y_{t_2}$ with $Y_i = f^{-1}(W_i)$ for $i = 1, \dots, t_2$. This proves Lemma 9. \square

We want to relax the condition of Lemma 9 that $\mathcal{O}_{B,S}^*/\mathcal{O}_S^*$ be finite and for this, we need some preparations.

We recall that a K -subalgebra B of A is said to be S -minimal if $1_A \in B$, and if B has no proper K -subalgebra B' with $1_A \in B'$ for which $\mathcal{O}_{B,S}^*/\mathcal{O}_{B',S}^*$ is finite. Every K -subalgebra B of A with $1_A \in B$ has an S -minimal K -subalgebra B' for which $\mathcal{O}_{B,S}^*/\mathcal{O}_{B',S}^*$ is finite. Namely, let B' be the intersection of all K -subalgebras B_1 of B with $1_A \in B_1$ for which $\mathcal{O}_{B,S}^*/\mathcal{O}_{B_1,S}^*$ is finite. Then $\mathcal{O}_{B',S}^*$ is the intersection of all groups $\mathcal{O}_{B_1,S}^*$. Furthermore, B has only finitely many K -subalgebras. Hence $\mathcal{O}_{B,S}^*/\mathcal{O}_{B',S}^*$ is finite. If B'' is a K -subalgebra of B' with $1_A \in B''$ such that $\mathcal{O}_{B',S}^*/\mathcal{O}_{B'',S}^*$ is finite, then $\mathcal{O}_{B,S}^*/\mathcal{O}_{B'',S}^*$ is finite, and therefore $B'' \supseteq B'$. Hence B' is S -minimal.

In what follows, let

$$B = \{x \in A : xV \subseteq V\}$$

be the algebra of scalars of A , and let B' be an S -minimal K -subalgebra of B for which $\mathcal{O}_{B,S}^*/\mathcal{O}_{B',S}^*$ is finite. Every K -subalgebra of A is semi-simple, i.e. isomorphic to a direct sum of finite extension fields of K . So in particular we have

$$B' \cong L'_1 \oplus \dots \oplus L'_q$$

for certain finite extension fields L'_1, \dots, L'_q of K . Then B' has K -subalgebras L''_1, \dots, L''_q such that

$$(4.5) \quad B' = L''_1 + \dots + L''_q \text{ as vector space, } L''_i \cdot L''_j = (0) \text{ for } 1 \leq i < j \leq q,$$

$$L''_i \cong L'_i \text{ for } i = 1, \dots, q.$$

For $i = 1, \dots, q$, denote by 1_i the unit element of L''_i . (4.5) and $1_A \in B'$ imply that

$$(4.6) \quad 1_A = 1_1 + \dots + 1_q, \quad 1_i \cdot 1_j = 0 \text{ for } 1 \leq i < j \leq q.$$

Let $1_i = (\xi_{i1}, \dots, \xi_{it})$ with $\xi_{ij} \in M_j$ for $j = 1, \dots, t$. Since $1_i^2 = 1_i$, we have $\xi_{ij}^2 = \xi_{ij}$, whence $\xi_{ij} \in \{0, 1\}$ for $j = 1, \dots, t$. Together with (4.6) this implies that there are subsets P_1, \dots, P_q of $\{1, \dots, t\}$ such that

$$(4.7) \quad 1_i = (\xi_{i1}, \dots, \xi_{it}) \text{ with } \xi_{ij} = 1 \text{ for } j \in P_i, \quad \xi_{ij} = 0 \text{ for } j \notin P_i,$$

$$(4.8) \quad P_1 \cup \dots \cup P_q = \{1, \dots, t\}, \quad P_i \cap P_j = \emptyset \text{ for } 1 \leq i < j \leq q.$$

Define the K -algebras

$$A_i = \bigoplus_{j \in P_i} M_j \text{ for } i = 1, \dots, q,$$

the projections

$$\Pi_i : A \mapsto A_i : (\xi_1, \dots, \xi_t) \mapsto (\xi_j : j \in P_i) \text{ for } i = 1, \dots, q,$$

and

$$\Pi = (\Pi_1, \dots, \Pi_q) : A \mapsto A_1 \oplus \dots \oplus A_q : x \mapsto (\Pi_1(x), \dots, \Pi_q(x)).$$

Π is merely a permutation of coordinates, so Π is a K -algebra isomorphism from A to $A_1 \oplus \dots \oplus A_q$. Further define

$$B_i := \Pi_i(B), \quad L_i := \Pi_i(B'), \quad V_i := \Pi_i(V) \text{ for } i = 1, \dots, q,$$

where B_i, L_i are K -subalgebras, and V_i is a subspace of A_i . Then we have:

Lemma 10. (i). $\Pi(B) = B_1 \oplus \dots \oplus B_q$, $\Pi(B') = L_1 \oplus \dots \oplus L_q$, $\Pi(V) = V_1 \oplus \dots \oplus V_q$.

(ii). For $i = 1, \dots, q$, L_i is isomorphic to a finite extension field of K .

(iii). $B_i = \{x \in A_i : xV_i \subseteq V_i\}$ for $i = 1, \dots, q$.

Proof. (i). We prove only that $\Pi(V) = V_1 \oplus \dots \oplus V_q$; the proofs that $\Pi(B) = B_1 \oplus \dots \oplus B_q$ and $\Pi(B') = L_1 \oplus \dots \oplus L_q$ are entirely similar. It is obvious that $\Pi(V) \subseteq V_1 \oplus \dots \oplus V_q$. Conversely, let $x = (x_1, \dots, x_q)$ with $x_j \in V_j$ for $j = 1, \dots, q$. Choose $y_j \in V$ such that $\Pi_j(y_j) = x_j$ for $j = 1, \dots, q$ and put $y := \sum_{j=1}^q 1_j \cdot y_j$. Since $1_j \in L_j'' \subseteq B' \subseteq B$ we have $1_j V \subseteq V$ for $j = 1, \dots, q$; hence $y \in V$. Now (4.7) and (4.8) imply that for $j = 1, \dots, q$, the coordinates of y with indices in P_j are equal to the corresponding coordinates of y_j . Hence $\Pi_j(y) = \Pi_j(y_j) = x_j$ for $j = 1, \dots, q$. Therefore, $\Pi(y) = x$. We infer that indeed $\Pi(V) = V_1 \oplus \dots \oplus V_q$.

(ii). Let $i \in \{1, \dots, q\}$. We first show that $\Pi_i(L_i'') = \Pi_i(B')$. L_i'' is a K -subalgebra of B' , hence $\Pi_i(L_i'') \subseteq \Pi_i(B')$. Conversely, let $x \in B'$. Then $x = x_1 + \dots + x_q$ with $x_j \in L_j''$ for $j = 1, \dots, q$. Now $\Pi_i(1_i) = (1, \dots, 1)$ and by (4.5) we have $1_i x_j = 0$ for $j \neq i$. Hence

$$\Pi_i(x) = \Pi_i(1_i x) = \Pi_i(1_i x_i) = \Pi_i(x_i) \in \Pi_i(L_i'').$$

This shows that indeed $\Pi_i(L_i'') = \Pi_i(B')$. Now Π_i is non-trivial as its image contains $(1, \dots, 1)$ and L_i'' is a field, hence $L_i = \Pi_i(L_i'')$ is a field.

(iii). Let $i \in \{1, \dots, q\}$. Put $\tilde{B}_i := \{x \in A_i : xV_i \subseteq V_i\}$. For $x \in B_i$ we have $x = \Pi_i(y)$ for some $y \in B$, whence $xV_i = \Pi_i(yV) \subseteq \Pi_i(V) = V_i$. Therefore, $B_i \subseteq \tilde{B}_i$. To prove the opposite inclusion, consider $\tilde{B} = \Pi^{-1}(\tilde{B}_1 \oplus \dots \oplus \tilde{B}_q)$. Then \tilde{B} is a K -subalgebra of A and for $x \in \tilde{B}$ we have by (i) $xV = \Pi^{-1}(\Pi(x) \cdot (V_1 \oplus \dots \oplus V_q)) \subseteq \Pi^{-1}(V_1 \oplus \dots \oplus V_q) = V$; therefore, $\tilde{B} \subseteq B$. It follows that $\tilde{B}_i \subseteq \Pi_i(\tilde{B}) \subseteq \Pi_i(B) = B_i$, which completes the proof. \square

Fix again $i \in \{1, \dots, q\}$. We have $L_i \subseteq B_i \subseteq A_i$, so that A_i may be viewed as an L_i -algebra and B_i as an L_i -subalgebra of A_i . Further, the unit element 1_{A_i} of A_i is just the unit element of L_i , and so $1_{A_i} \in B_i$. Lastly, by (iii) of Lemma

10, V_i is an L_i -vector space. Note that $\mathcal{O}_{A_i,S} = \bigoplus_{j \in P_i} \mathcal{O}_{M_j,S}$, $\mathcal{O}_{B_i,S} = \mathcal{O}_{A_i,S} \cap B_i$, $\mathcal{O}_{L_i,S} = \mathcal{O}_{A_i,S} \cap L_i$ are the integral closures of \mathcal{O}_S in A_i, B_i, L_i , respectively. Clearly, $\mathcal{O}_{B_i,S}^*/\mathcal{O}_{L_i,S}^*$ is a homomorphic image of $\mathcal{O}_{B,S}^*/\mathcal{O}_{B',S}^*$, so

$$(4.9) \quad \mathcal{O}_{B_i,S}^*/\mathcal{O}_{L_i,S}^* \text{ is finite.}$$

We are now ready to prove the following generalisation of Lemma 9:

Lemma 11. *We have that either $V = yB'$ for some $y \in A$, or there are proper K -linear subspaces Y_1, \dots, Y_{t_3} of V such that*

$$V \cap x_0 \mathcal{O}_{A,S}^* \subseteq Y_1 \cup \dots \cup Y_{t_3} \text{ with } t_3 \leq (2^{66} r^4)^{n^2 s}.$$

Proof. As mentioned before, for $i = 1, \dots, q$, V_i may be viewed as an L_i -vector space. First assume that $\dim_{L_i} V_i = 1$ for $i = 1, \dots, q$. Then for $i = 1, \dots, q$ there is an $y_i \in A_i$, such that $V_i = y_i L_i$. Together with part (i) of Lemma 10 this implies that $V = \Pi^{-1}(y_1 L_1 \oplus \dots \oplus y_q L_q) = yB'$ with $y = \Pi^{-1}((y_1, \dots, y_q))$.

Now assume that $\dim_{L_1} V_1 \geq 2$, say. Put $n_1 := \dim_{L_1} V_1$, $r_1 := \dim_{L_1} A_1$, let S_1 be the set of places lying above those in S , and s_1 the cardinality of S_1 . Then since V_1 is a K -linear subspace of $\Pi(V) \cong V$, and A_1 of $\Pi(A) \cong A$, we have

$$n_1[L_1 : K] = \dim_K V_1 \leq n, \quad r_1[L_1 : K] = \dim_K A_1 \leq r,$$

$$s_1 \leq s[L_1 : K].$$

Further, putting $x'_0 := \Pi_1(x_0)$, we have

$$\Pi_1(V \cap x_0 \mathcal{O}_{A,S}^*) \subseteq V_1 \cap x'_0 \mathcal{O}_{A,S}^*.$$

In view of part (iii) of Lemma 10 and of (4.9), we may apply Lemma 9 with L_1, A_1, B_1, V_1, S_1 replacing K, A, B, V, S . Thus, there are proper L_1 -linear subspaces Z_1, \dots, Z_{t_3} of V_1 , with

$$t_3 \leq (2^{66} r_1^4)^{n_1^2 s_1} \leq (2^{66} r^4)^{n^2 s}$$

such that $V_1 \cap x'_0 \mathcal{O}_{A_1,S}^* \subseteq Z_1 \cup \dots \cup Z_{t_3}$. But each of these subspaces Z_j is a K -linear subspace of V_1 . Hence it follows that $V \cap x_0 \mathcal{O}_{A,S}^* \subseteq Y_1 \cup \dots \cup Y_{t_3}$ where $Y_j = \Pi_1^{-1}(Z_j)$ is a proper K -linear subspace of V . This proves Lemma 11. \square

We recall that $e(n)$ is defined by $e(n) = \frac{1}{3}n(n+1)(2n+1) - 2$.

Lemma 12. *There are $y_1, \dots, y_{t_4} \in A^*$ and S -minimal K -subalgebras B_1, \dots, B_{t_4} of A , such that*

$$y_i B_i \subseteq V \text{ for } i = 1, \dots, t_4,$$

$$V \cap x_0 \mathcal{O}_{A,S}^* \subseteq y_1 \mathcal{O}_{B_1,S}^* \cup \dots \cup y_{t_4} \mathcal{O}_{B_{t_4},S}^* \text{ with } t_4 \leq (2^{33} r^2)^{e(n)s}.$$

Proof. We first deal with the special case that $V = yB_1$ for some $y \in A$ and some S -minimal K -subalgebra B_1 of A . Assume that $V \cap x_0\mathcal{O}_{A,S}^* \neq \emptyset$ and let $y_1 \in V \cap x_0\mathcal{O}_{A,S}^*$. Then $x_0\mathcal{O}_{A,S}^* = y_1\mathcal{O}_{A,S}^*$. By assumption we have $x_0 \in A^*$, hence it follows that $y_1 \in A^*$. Further, $y_1 = yz$ for some $z \in B_1$, and so $z \in B_1^*$. Therefore, $V = yB_1 = y_1B_1$. It follows that

$$V \cap x_0\mathcal{O}_{A,S}^* = y_1B_1 \cap y_1\mathcal{O}_{A,S}^* = y_1\mathcal{O}_{B_1,S}^*,$$

which implies Lemma 12 for $V = yB_1$.

We prove Lemma 12 in full generality by induction on $n = \dim_K V$. If $n = 1$, then $V = yK$ for some $y \in A$ and we are done since K is an S -minimal subalgebra of A . Suppose that $n \geq 2$, and that V is not equal to yB for some $y \in A$ and some S -minimal K -subalgebra B of A . Then by Lemma 11 we have $V \cap x_0\mathcal{O}_{A,S}^* \subseteq Y_1 \cup \dots \cup Y_{t_3}$ with $t_3 \leq (2^{66}r^4)^{n^2s}$, where Y_1, \dots, Y_{t_3} are proper K -linear subspaces of V . Now by the induction hypothesis we have for $i = 1, \dots, t_3$,

$$Y_i \cap x_0\mathcal{O}_{A,S}^* \subseteq y_{i,1}\mathcal{O}_{B_{i,1},S}^* \cup \dots \cup y_{i,t_5}\mathcal{O}_{B_{i,t_5},S}^* \text{ with } t_5 \leq (2^{33}r^2)^{e(n-1)s}$$

where $y_{i,j} \in A^*$, and $B_{i,j}$ is an S -minimal K -subalgebra of A with $y_{i,j}B_{i,j} \subseteq Y_i$ for $j = 1, \dots, t_5$. It follows that

$$V \cap x_0\mathcal{O}_{A,S}^* \subseteq \bigcup_{i=1}^{t_3} \bigcup_{j=1}^{t_5} y_{i,j}\mathcal{O}_{B_{i,j},S}^* \text{ with } y_{i,j}B_{i,j} \subseteq V.$$

Since $t_3t_5 \leq (2^{33}r^2)^{\{2n^2+e(n-1)\}s} = (2^{33}r^2)^{e(n)s}$, this proves Lemma 12. \square

Before finishing the proof of Theorem 1, we prove the following Lemma:

Lemma 13. *Let B be an S -minimal K -subalgebra of A , and $x_0\mathfrak{U}_{\mathcal{M},B}$ an (\mathcal{M}, B) -family of solutions of (1.7), with $x_0 \in \mathcal{M}^B$. Then $x_0\mathfrak{U}_{\mathcal{M},B}$ is irreducible.*

Proof. Suppose that $x_0\mathfrak{U}_{\mathcal{M},B}$ is reducible. Then there are proper subfamilies $x_1\mathfrak{U}_{\mathcal{M},B_1}, \dots, x_w\mathfrak{U}_{\mathcal{M},B_w}$ of $x_0\mathfrak{U}_{\mathcal{M},B}$ such that

$$(4.10) \quad x_0\mathfrak{U}_{\mathcal{M},B} = x_1\mathfrak{U}_{\mathcal{M},B_1} \cup \dots \cup x_w\mathfrak{U}_{\mathcal{M},B_w}.$$

Further, there is no loss of generality to assume that

$$(4.11) \quad x_i \in \mathcal{M}^{B_i}, \quad B_i \subsetneq B \text{ for } i = 1, \dots, w.$$

Namely, if for instance B_1 is not a K -subalgebra of B then by Lemma 3, $x_1\mathfrak{U}_{\mathcal{M},B_1} = x\mathfrak{U}_{\mathcal{M},B} \cap x_1\mathfrak{U}_{\mathcal{M},B_1}$ is the union of finitely many $(\mathcal{M}, B \cap B_1)$ -families and, in (4.10), we may replace $x_1\mathfrak{U}_{\mathcal{M},B_1}$ by this union. Further, if $B_1 = B$ then $x_1\mathfrak{U}_{\mathcal{M},B_1}$ is not a proper subfamily of $x_0\mathfrak{U}_{\mathcal{M},B}$.

Put $\rho_B := \text{rank } \mathcal{O}_{B,S}^*/\mathcal{O}_S^*$, $\rho := \max_{i=1, \dots, w} \{\text{rank } \mathcal{O}_{B_i,S}^*/\mathcal{O}_S^*\}$. From (4.11) and the fact that B is S -minimal, it follows that $\rho < \rho_B$. On the other hand, letting

$N_{\mathcal{F}}(X)$ be the quantity in the statement of Lemma 2, it follows from Lemma 2 and (4.10) that

$$N_{x_0\mathfrak{U}_{\mathcal{M},B}}(X) = \gamma(\log X)^{\rho_B} + O((\log X)^{\rho_B-1}) \text{ as } X \rightarrow \infty \text{ with } \gamma > 0,$$

$$N_{x_0\mathfrak{U}_{\mathcal{M},B}}(X) = N_{\cup_{i=1}^w x_i\mathfrak{U}_{\mathcal{M},B_i}}(X) = O((\log X)^{\rho}) \text{ as } X \rightarrow \infty.$$

Thus, the assumption that $x_0\mathfrak{U}_{\mathcal{M},B}$ is reducible leads to a contradiction. This proves Lemma 13. \square

Proof of Theorem 1. By Lemma 14, the set of solutions of (1.7) is contained in some union $\cup_{j=1}^{t_1} \{V \cap x_j\mathcal{O}_{A,S}^*\}$ with $x_j \in A^*$ for $j = 1, \dots, t_1$ and $t_1 \leq \psi_2(\beta)$. By Lemma 12, for $j = 1, \dots, t_1$, $V \cap x_j\mathcal{O}_{A,S}^*$ is a subset of some finite union $\cup_{h=1}^{t_{4j}} y_{jh}\mathcal{O}_{B_{jh},S}^*$ with $t_{4j} \leq (2^{33}r^2)^{e(n)s}$, where $y_{jh} \in A^*$ and B_{jh} is an S -minimal K -subalgebra of A with $y_{jh}B_{jh} \subseteq V$, $h = 1, \dots, t_{4j}$. It follows that the set of solutions of (1.7) is contained in $\cup_{h=1}^w y_h\mathcal{O}_{B_h,S}^*$ with $w \leq (2^{33}r^2)^{e(n)s} \psi_2(\beta)$, where $y_h \in A^*$ and B_h is an S -minimal K -subalgebra of A with $y_hB_h \subseteq V$, $h = 1, \dots, w$.

We recall that if B is an S -minimal K -subalgebra of A , then, by Lemma 13, any (\mathcal{M}, B) -family of solutions is automatically irreducible. Hence the proof of Theorem 1 is complete once we have shown that the set of solutions of (1.7) belonging to some coset $y\mathcal{O}_{B,S}^*$ with $y \in A^*$, $yB \subseteq V$ is the union of at most $I := [\mathcal{O}_{B,S}^* : \mathfrak{U}_{\mathcal{M},B}]$ (\mathcal{M}, B) -families of solutions. Clearly, $y\mathcal{O}_{B,S}^*$ is the union of I cosets $z\mathfrak{U}_{\mathcal{M},B}$ with $z \in A^*$. Suppose that $z\mathfrak{U}_{\mathcal{M},B}$ contains a solution, say z_0 , of (1.7). Then $z\mathfrak{U}_{\mathcal{M},B} = z_0\mathfrak{U}_{\mathcal{M},B}$. We have $z_0 \in \mathcal{M}$ and also $z_0B = zB = yB \subseteq V$, so $z_0 \in V^B \cap \mathcal{M} = \mathcal{M}^B$, which implies that $z \in \mathcal{M}^B$. This proves that $z\mathfrak{U}_{\mathcal{M},B}$ is an (\mathcal{M}, B) -family of solutions of (1.7). This completes the proof of Theorem 1. \square

5. PROOF OF THEOREM 2

We will prove Theorem 2 more generally, for arbitrary fields K of characteristic 0. Thus, let K be any field of characteristic 0, $A = M_1 \oplus \dots \oplus M_t$ where M_1, \dots, M_t are finite extension fields of K with $\dim_K A = \sum_{i=1}^t [M_i : K] = r$, and V is an n -dimensional K -linear subspace of A . It is our purpose to prove that there are at most $\{n \max(r-n, 2)\}^n$ K -subalgebras of A with

$$(1.16) \quad 1_A \in B, \quad V^B \cap A^* \neq \emptyset.$$

We make some reductions. Let \overline{K} be the algebraic closure of K and $\overline{A} = \overline{K}^r$ with coordinatewise addition and multiplication. For $x = (\xi_1, \dots, \xi_t) \in A$, put $\mathfrak{f}(x) := (\tau_{1,1}(\xi_1), \dots, \tau_{1,r_1}(\xi_1), \dots, \tau_{t,1}(\xi_t), \dots, \tau_{t,r_t}(\xi_t))$, where for $i = 1, \dots, t$ $\tau_{i,1}, \dots, \tau_{i,r_i}$ ($r_i = [M_i : K]$) are the K -isomorphic embeddings of M_i into \overline{K} . Then \mathfrak{f} is an injective K -algebra homomorphism from A into \overline{A} . It is easy to check that \mathfrak{f} maps K -linearly independent elements of A to \overline{K} -linearly independent elements of \overline{A} . Hence, if for a K -linear subspace W of A we define \overline{W} to be the \overline{K} -vector space generated by

$f(W)$, we have that $\dim_{\overline{K}} \overline{W} = \dim_K W$ and that W is uniquely determined by \overline{W} . Finally, if B is a K -subalgebra of A then \overline{B} is a \overline{K} -subalgebra of \overline{A} : namely, if $x, y \in \overline{B}$, then $x = \sum \xi_i f(x_i)$, $y = \sum \eta_j f(y_j)$ with $\xi_i, \eta_j \in \overline{K}$, $x_i, y_j \in B$ and therefore, $xy = \sum \xi_i \eta_j f(x_i y_j) \in \overline{B}$. Note that $\mathbf{1} = (1, \dots, 1)$ (r times) is the element of A and that $\overline{A}^* = \{(\xi_1, \dots, \xi_r) \in \overline{K}^r : \xi_1 \dots \xi_r \neq 0\}$. For K -subalgebras B of A with (1.16) we have

$$(5.1) \quad \mathbf{1} \in \overline{B}, \quad \overline{V}^{\overline{B}} \cap \overline{A}^* \neq \emptyset.$$

Namely, it is clear that $\mathbf{1} \in \overline{B}$. Further, if $x \in V^B \cap A^*$, we have $f(x) \in \overline{A}^*$ and also $xB \subseteq V$, whence $f(x)\overline{B} \subseteq \overline{V}$, i.e. $f(x) \in \overline{V}^{\overline{B}} \cap \overline{A}^*$. Since B is uniquely determined by \overline{B} , it follows that the number of K -subalgebras B of A with (1.16) is at most the number of \overline{K} -subalgebras \overline{B} of \overline{A} with (5.1). Hence it suffices to prove the following:

Lemma 14. *\overline{A} has at most $\{n \max(r - n, 2)\}^n$ \overline{K} -subalgebras \overline{B} with (5.1).*

Proof. Let \overline{B} be a \overline{K} -subalgebra of \overline{A} with (5.1). Then, for some $q \leq r$, \overline{B} is isomorphic to \overline{K}^q with coordinatewise operations. This implies that \overline{B} has \overline{K} -subalgebras L''_1, \dots, L''_q such that $L''_i \cong \overline{K}$ for $i = 1, \dots, q$, $L''_1 + \dots + L''_q = \overline{B}$, and $L''_i \cdot L''_q = (0)$ for $1 \leq i < j \leq q$. Letting 1_i be the unit element of L''_i for $i = 1, \dots, q$, we find, completely similarly to (4.7) and (4.8), that there are non-empty subsets P_1, \dots, P_q of $\{1, \dots, r\}$ such that

$$(5.2) \quad 1_i = (\xi_{i1}, \dots, \xi_{ir}) \text{ with } \xi_{ij} = 1 \text{ for } j \in P_i, \xi_{ij} = 0 \text{ for } j \notin P_i,$$

$$(5.3) \quad P_1 \cup \dots \cup P_q = \{1, \dots, r\}, \quad P_i \cap P_j = \emptyset \text{ for } 1 \leq i < j \leq r.$$

First suppose that $r > n$. On noting that $\dim_{\overline{K}} \overline{V} = n$, after a permutation of coordinates if necessary, we may assume that \overline{V} is the set of solutions (ξ_1, \dots, ξ_r) of a system of linear equations

$$(5.4) \quad \xi_k = \sum_{j=1}^n c_{kj} \xi_j \text{ for } k = n+1, \dots, r,$$

with $c_{kj} \in \overline{K}$. Let $(\xi_1, \dots, \xi_r) \in \overline{V}^{\overline{B}} \cap \overline{A}^*$. Then $1_i x \in \overline{V}$ for $i = 1, \dots, q$. (5.2) implies that the coordinates of $1_i x$ with indices in P_i are the same as those of x , while the coordinates of $1_i x$ with indices outside P_i are 0. Together with (5.4) this implies

$$(5.5) \quad \begin{cases} \xi_k = \sum_{j \in Q_i} c_{kj} \xi_j & \text{for } k \in R_i, \quad i = 1, \dots, q, \\ 0 = \sum_{j \in Q_i} c_{kj} \xi_j & \text{for } k \in \tilde{R}_i := \{n+1, \dots, r\} \setminus R_i, \quad i = 1, \dots, q, \end{cases}$$

where $Q_i := P_i \cap \{1, \dots, n\}$, $R_i := P_i \cap \{n+1, \dots, r\}$, $i = 1, \dots, q$. Note that

$$(5.6) \quad \begin{cases} Q_1 \cup \dots \cup Q_q = \{1, \dots, n\}, & Q_i \cap Q_j = \emptyset & \text{for } 1 \leq i < j \leq q, \\ R_1 \cup \dots \cup R_q = \{n+1, \dots, r\}, & R_i \cap R_j = \emptyset & \text{for } 1 \leq i < j \leq q, \\ Q_i \cap R_j \neq \emptyset & \text{for } i, j = 1, \dots, q. \end{cases}$$

Further, by (5.2) and the fact that $\overline{B} = L_1'' + \dots + L_q'' = 1_1 \overline{K} + \dots + 1_q \overline{K}$, we have that \overline{B} is determined uniquely by P_1, \dots, P_q , whence by $Q_1, \dots, Q_q, R_1, \dots, R_q$. Recalling that $x \in A^*$ we infer that it suffices to prove

$$(5.7) \quad \begin{aligned} & \text{there are at most } \{n \max(r-n, 2)\}^n \text{ collections } \{Q_1, \dots, Q_q, R_1, \dots, R_q\} \\ & \text{with (5.6) such that (5.5) has a solution with } \xi_1 \dots \xi_r \neq 0. \end{aligned}$$

For the moment, we fix Q_1, \dots, Q_q and determine an upper bound for the number of collections $\{R_1, \dots, R_q\}$ for which (5.5) has a solution with $\xi_1 \dots \xi_r \neq 0$. Let $n_i := \#Q_i$ for $i = 1, \dots, q$. Take $i \in \{1, \dots, q\}$. We have $Q_i \neq \emptyset$ since otherwise $R_i \neq \emptyset$ and each solution of (5.5) has $\xi_k = 0$ for $k \in R_i$. Define the vectors $\mathbf{c}_k = (c_{kj} : j \in Q_i)$ ($k = n+1, \dots, r$). We have $\text{rank}\{\mathbf{c}_k : k \in \tilde{R}_i\} \leq n_i - 1$, since otherwise each solution of (5.5) has $\xi_j = 0$ for $j \in Q_i$. Further, for each $l \in R_i$ the vector \mathbf{c}_l is linearly independent of $\{\mathbf{c}_k : k \in \tilde{R}_i\}$, since otherwise the equations $\sum_{j \in Q_i} c_{kj} \xi_j = 0$ for $k \in \tilde{R}_i$ imply $\sum_{j \in Q_i} c_{lj} \xi_j = 0$ for some $l \in R_i$ and so each solution of (5.5) has $\xi_l = 0$. It follows that $\{\mathbf{c}_k : k \in \tilde{R}_i\}$ consists of all vectors in $\{\mathbf{c}_k : k = n+1, \dots, r\}$ that are linear combinations of some linearly independent subset of $\{\mathbf{c}_k : k \in \tilde{R}_i\}$. But then, this linearly independent subset uniquely determines R_i . Recalling that $\text{rank}\{\mathbf{c}_k : k \in \tilde{R}_i\} \leq n_i - 1$, we infer that the number of possibilities for R_i is at most the number of linearly independent subsets of $\{\mathbf{c}_k : k = n+1, \dots, r\}$ of cardinality $\leq n_i - 1$, and the latter is at most

$$\binom{r-n}{0} + \binom{r-n}{1} + \dots + \binom{r-n}{n_i-1} \leq \{\max(r-n, 2)\}^{n_i}.$$

Therefore, for given Q_1, \dots, Q_q , the number of possibilities for $\{R_1, \dots, R_q\}$ is at most

$$\{\max(r-n, 2)\}^{n_1 + \dots + n_q} = \{\max(r-n, 2)\}^n.$$

The number of possibilities for $\{Q_1, \dots, Q_q\}$ is at most the number of partitions of $\{1, \dots, n\}$ into disjoint sets which is $\leq n^n$. This implies (5.7), hence Lemma 14 for $r > n$. If $r = n$, then the sets R_1, \dots, R_q do not occur and we have only to estimate the number of possibilities for $\{Q_1, \dots, Q_q\}$. So in that case, Lemma 14 follows also. \square

REFERENCES

- [1] Z.I. Borevich, I.R. Shafarevich, *Number Theory*, Academic Press, New York, 1967.
- [2] G.R. Everest, *On the solution of the norm form equation*, Amer. J. Math. **114** (1992), 667-681, Addendum, *ibid*, 787-788.

- [3] G.R. Everest, K. Györy, *Counting solutions of decomposable form equations*, Acta Arith. (to appear).
- [4] J.-H. Evertse, *The number of solutions of decomposable form equations*, Invent. math. **122** (1995), 559-601.
- [5] K. Györy, *On the numbers of families of solutions of systems of decomposable form equations*, Publ. Math. Debrecen **42** (1993), 65-101.
- [6] K. Györy, A. Pethő, *Über der Verteilung der Lösungen von Normformen Gleichungen II*, Acta Arith. **32** (1977), 349-363.
- [7] K. Györy, A. Pethő, *Über der Verteilung der Lösungen von Normformen Gleichungen III*, Acta Arith. **37** (1980), 143-165.
- [8] S. Lang, *Fundamentals of Diophantine Geometry*, Springer Verlag, Berlin, etc., 1983..
- [9] M. Laurent, *Equations diophantiennes exponentielles*, Invent. math. **78** (1984), 299-327.
- [10] D.G. Northcott, *An inequality in the theory of arithmetic on algebraic varieties*, Proc. Cambridge Philos. Soc. **45** (1949), 502-509.
- [11] D.G. Northcott, *A further inequality in the theory of arithmetic on algebraic varieties*, Proc. Cambridge Philos. Soc. **45** (1949), 510-518.
- [12] A. Pethő, *Über die Verteilung der Lösungen von S-Normformen Gleichungen*, Publ. Math. Debrecen **29** (1982), 1-17.
- [13] H.P. Schlickewei, *On norm form equations*, J. Number Th. **9** (1977), 370-380.
- [14] H.P. Schlickewei, *S-unit equations over number fields*, Invent. math. **102** (1990), 95-107.
- [15] W.M. Schmidt, *Linearformen mit algebraischen Koeffizienten II*, Math. Ann. **191** (1971), 1-20.
- [16] W.M. Schmidt, *Norm form equations*, Ann. Math. **96** (1972), 525-551.
- [17] W.M. Schmidt, *The number of solutions of norm form equations*, Trans. Am. Math. Soc. **317** (1990), 197-227.
- [18] P.M. Voutier, *Effective and quantitative results on integral solutions of certain classes of Diophantine equations*, Ph.D.-thesis, University of Colorado at Boulder, 1993.

Adresses of the authors:

University of Leiden, Mathematical Institute,
 P.O. Box 9512, 2300 RA Leiden, The Netherlands,
 e-mail evertse@wi.leidenuniv.nl

Kossuth Lajos University, Mathematical Institute,
 4010 Debrecen, P.O. Box 12, Hungary,
 e-mail gyory@math.klte.hu