Answer the following questions, ready for the test on thursday.

In Exercises 1 and 2, compute each matrix sum or product if it is defined. If an expression is undefined, explain why. Let

$$A = \begin{bmatrix} 2 & 0 & -1 \\ 4 & -5 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 7 & -5 & 1 \\ 1 & -4 & -3 \end{bmatrix},$$
$$C = \begin{bmatrix} 1 & 2 \\ -2 & 1 \end{bmatrix}, \quad D = \begin{bmatrix} 3 & 5 \\ -1 & 4 \end{bmatrix}, \quad E = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$$
$$1. \quad -2A, \quad B - 2A, \quad AC, \quad CD$$
$$2. \quad A + 3B, \quad 2C - 3E, \quad DB, \quad EC$$

In Exercises 5 and 6, compute the product AB in two ways: (a) by the definition, where  $A\mathbf{b}_1$  and  $A\mathbf{b}_2$  are computed separately, and (b) by the row–column rule for computing AB.

**5.** 
$$A = \begin{bmatrix} -1 & 3 \\ 2 & 4 \\ 5 & -3 \end{bmatrix}, B = \begin{bmatrix} 4 & -2 \\ -2 & 3 \end{bmatrix}$$

- If a matrix A is 5 × 3 and the product AB is 5 × 7, what is the size of B?
- 8. How many rows does B have if BC is a  $5 \times 4$  matrix?

9. Let 
$$A = \begin{bmatrix} 2 & 3 \\ -1 & 1 \end{bmatrix}$$
 and  $B = \begin{bmatrix} 1 & 9 \\ -3 & k \end{bmatrix}$ . What value(s) of k, if any, will make  $AB = BA$ ?

- **11.** Let  $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ 3 & 5 & 6 \end{bmatrix}$  and  $D = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ . Compute *AD* and *DA*. Explain how the columns or rows of *A* change when *A* is multiplied by *D* on the right or on the left. Find a 3 × 3 matrix *B*, not the identity matrix or the zero matrix, such that AB = BA.
- **12.** Let  $A = \begin{bmatrix} 3 & -6 \\ -2 & 4 \end{bmatrix}$ . Construct a 2 × 2 matrix *B* such that *AB* is the zero matrix. Use two different nonzero columns for *B*.

Exercises 15 and 16 concern arbitrary matrices A, B, and C for which the indicated sums and products are defined. Mark each statement True or False. Justify each answer.

- **15.** a. If A and B are  $2 \times 2$  matrices with columns  $\mathbf{a}_1$ ,  $\mathbf{a}_2$ , and  $\mathbf{b}_1$ ,  $\mathbf{b}_2$ , respectively, then  $AB = [\mathbf{a}_1\mathbf{b}_1 \quad \mathbf{a}_2\mathbf{b}_2]$ .
  - b. Each column of AB is a linear combination of the columns of B using weights from the corresponding column of A.
  - c. AB + AC = A(B + C)
  - d.  $A^T + B^T = (A + B)^T$
  - e. The transpose of a product of matrices equals the product of their transposes in the same order.

- **31.** Show that  $I_m A = A$  where A is an  $m \times n$  matrix. Assume  $I_m \mathbf{x} = \mathbf{x}$  for all  $\mathbf{x}$  in  $\mathbb{R}^m$ .
- **32.** Show that  $AI_n = A$  when A is an  $m \times n$  matrix. [*Hint:* Use the (column) definition of  $AI_n$ .]