

Introduction to index theory

Yuezhao Li*

Mathematical Institute, Leiden University

11 November, 2021

This is a handout for a reading seminar at Leiden University. In this talk, I will give a very basic introduction on the celebrated Atiyah–Singer index theorem, which is one of the most exciting and elegant mathematical achievements in the past century. The theorem claims that the analytical index of an elliptic (pseudo)differential operator, which is the Fredholm index of this operator (viewed as a linear operator between suitable Hilbert spaces), equals the topological index of the operator, defined in purely topological terms. I will define the analytical index and the topological index of an elliptic operator, and sketch the proof why they are equal. The proof will be closely following the original K-theoretic proof due to Atiyah and Singer [AS1], but I will also try to highlight the role of K-theory of C*-algebras.

Notations and Conventions

- Let \mathcal{H} be a complex Hilbert space. We write $\mathbb{K}(\mathcal{H})$ for the C*-algebra of compact operators on \mathcal{H} , and write $\mathbb{B}(\mathcal{H})$ for the C*-algebra of bounded operators on \mathcal{H} . When the Hilbert space is clear from the context we shall also omit \mathcal{H} and write \mathbb{K} and \mathbb{B} for simplicity.
- We write \hookrightarrow for monomorphisms and \twoheadrightarrow for epimorphisms.
- We write H_c for the compactly-supported cohomology, **with coefficients in \mathbb{Q}** . We write H_c^{even} for the direct sum of all cohomology groups in even dimensions. H_c^{odd} is defined likewise.
- We write K_0, K_1, \dots for K-theory of C*-algebras, and K^0, K^{-1}, \dots for **compactly-supported** K-theory of topological spaces. For simplicity (and in consistency with literature) we will usually use K instead of K^0 . For instance, let X be a locally compact topological space. Then we have

$$K_0(C_0(X)) = K^0(X) = K(X).$$

- Let $E \rightarrow X$ be a complex vector bundle of complex rank k . Then it is both oriented and K-oriented (Proposition 3.1). We write $\phi_E \in K(E)$ for the Thom class of E in K-theory, and $\psi_E \in H_c^{\text{even}}(E)$ for the Thom class of E in compactly-supported cohomology.
- We use \smile to denote the cup product in (generalised) cohomology theories, like (compactly-supported) cohomology and K-theory. One can find their definitions in some standard references, for example, [Hat02, Hat03].

* <mailto:y.li@math.leidenuniv.nl>

Contents

1 An overview	2
2 The analytical index	3
2.1 K-theory and Fredholm index	3
2.2 Pseudodifferential extension	4
2.3 The $K(TM)$ -symbol class	6
3 The topological index	7
3.1 Thom isomorphism	7
3.2 The cohomological formula	8
4 Remarks on the general cases	10
5 Proof of the index theorem	10

1 An overview

The general form of an index theorem is given by an equation

$$\boxed{\text{Analytical index} = \text{Topological index}}$$

In the Atiyah–Singer index theorem, the **analytical index** is given by the Fredholm index of an elliptic (pseudo)differential operator. The **topological index** is given by some purely “topological” data coming from the symbol of the operator, together with the base manifold.



Theorem 1.1 (Atiyah–Singer)

Let D be an elliptic (pseudo)differential operator on a compact m -dimensional manifold M . Then

$$\text{a-ind}(D) = \text{t-ind}(D),$$

where

$$\text{a-ind} := \text{Index}(D),$$

and

$$\text{t-ind}(D) := (-1)^m \int_{TM} \pi^* \text{Td}(TM \otimes \mathbb{C}) \smile \text{Ch}([\sigma(D)]).$$

Atiyah–Singer index theorem vastly generalises the following “classical” theorems:

- Chern–Gauß–Bonnet theorem
- Hirzebruch–Riemann–Roch theorem.
- Hirzebruch signature theorem.

Before going into the details, one might wonder why such a theorem could exist. There are various discussions on this specific question, aiming at an insight of such a tremendous theorem. One superficial idea might be that the Fredholm index is homotopy-invariant:



Proposition 1.2

Let \mathcal{H} be a complex infinite-dimensional Hilbert space, $\mathcal{F}(\mathcal{H})$ be the set of bounded Fredholm operators on \mathcal{H} , equipped with the norm topology. Then the map

$$\mathcal{F}(\mathcal{H}) \rightarrow \mathbb{Z}, \quad T \mapsto \text{Index}(T)$$

is continuous. Therefore, the Fredholm index is locally constant.

The fact at least implies that the analytical index is actually “topological”, and one could expect a topological formula for it. A generalisation of the previous proposition is:



Theorem 1.3 (Atiyah–Jänich)

Let X be a compact Hausdorff space. Then

$$[X, \mathcal{F}(\mathcal{H})] \cong K(X).$$

where $[X, Y]$ denotes the set of homotopy classes of maps from X to Y .

The previous proposition is the special case of Atiyah–Jänich Theorem where $X = \text{pt}$. The theorem also motivates that we can define an index for “family of operators parametrised by X ” whereas it should take value in $K(X)$ instead of $K(\text{pt}) = \mathbb{Z}$. This is the starting point of [AS4].

2 The analytical index

From now on we shall always let D be a **scalar-valued elliptic pseudodifferential operator of order 0** on a **compact** m -dimensional manifold M , whose meaning will be explained soon. In Section 4 we shall explain how to generalise to more general cases.

2.1 K-theory and Fredholm index

The analytical index $\text{a-ind}(D)$ of D is just its Fredholm index (view D as a Fredholm operator on $L^2(M)$). The best way to understand the Fredholm indices and operators might be introducing the (topological) K-theory of C^* -algebras. Nevertheless we will only treat it as a blackbox. It suffices to know that:

1. Topological K-theory is a homology theory of C^* -algebras in the sense that: let

$$I \xrightarrow{i} E \xrightarrow{q} Q$$

be an extension of C^* -algebras (that is, I is a closed ideal in the C^* -algebra E). Then there is an induced long exact sequence

$$\dots \rightarrow K_i(I) \xrightarrow{i_*} K_i(E) \xrightarrow{q_*} K_i(Q) \xrightarrow{\partial} K_{i-1}(I) \rightarrow \dots$$

of abelian groups.

2. The biggest distinction between topological K-theory and algebraic K-theory (of rings) is that in topological K-theory we have **Bott periodicity**

$$K_{i+2}(A) \cong K_i(A)$$

for any C^* -algebra A . Combined with 1 we have a 6-term cyclic exact sequence

$$\begin{array}{ccccc} K_0(I) & \longrightarrow & K_0(E) & \longrightarrow & K_0(Q) \\ \uparrow & & & & \downarrow \\ K_1(Q) & \longleftarrow & K_1(E) & \longleftarrow & K_1(I). \end{array}$$

3. (Relation to topological K-theory of topological spaces). Let X be a locally compact topological space. Then $C_0(X)$ is a C^* -algebra, and $K_i(C_0(X)) \cong K^{-i}(X)$. The right-hand side is defined by $K^{-i}(X) := K(X \times \mathbb{R}^i)$, the (compactly supported) K-theory of $X \times \mathbb{R}^i$, introduced in Bram’s talk.

Now we are at the right place to define the analytical index. Recall that (see, e.g. [Ols94, Chapter 14]):



Definition 2.1

Let \mathcal{H} be a complex Hilbert space. An operator $T \in \mathbb{B}(\mathcal{H})$ is called **Fredholm** if it has closed image, has finite-dimensional kernel and finite-dimensional cokernel^a. The **Fredholm index** of T , denoted by $\text{Index}(T)$, is defined by

$$\text{Index}(T) := \dim \ker T - \dim \text{coker } T.$$

^aThe image being closed is actually an unnecessary condition: it is automatically true if T has finite-dimensional kernel and cokernel.

and



Theorem 2.2 (Atkinson)

$T \in \mathbb{B}(\mathcal{H})$ is Fredholm, if and only if it is invertible modulo compact operators, if and only if the image of T in $\mathbb{B}(\mathcal{H})/\mathbb{K}(\mathcal{H})$ is invertible.

We have an extension of C^* -algebras

$$\mathbb{K} \hookrightarrow \mathbb{B} \rightarrow \mathbb{B}/\mathbb{K}.$$

An invertible element in a C^* -algebra A defines a class in $K_1(A)$. If T is Fredholm, then its image in \mathbb{B}/\mathbb{K} is invertible and represents a class $[T]$ in $K_1(\mathbb{B}/\mathbb{K})$. The long exact sequence in K -theory yields a boundary map

$$\partial: K_1(\mathbb{B}/\mathbb{K}) \rightarrow K_0(\mathbb{K}) \cong \mathbb{Z}.$$

We have the following well-known result (see, e.g. [Ols94, Chapter 14]):



Theorem 2.3

The boundary map $K_1(\mathbb{B}/\mathbb{K}) \xrightarrow{\partial} K_0(\mathbb{K}) \cong \mathbb{Z}$ sends the class $[T] \in K_1(\mathbb{B}/\mathbb{K})$ to the Fredholm index $\text{Index}(T)$ of T .

2.2 Pseudodifferential extension

Pseudodifferential calculus is a very deep and complicated topic. We shall be very brief and ad hoc here.

Recall from Yufan’s talk that we can construct operators on $L^2(\mathbb{R}^n)$ by “quantising” (that is, applying inverse Fourier transforms) “nice” functions

$$a(x, \xi) \mapsto A = \text{Op}(a) = a(x, -i\partial_x),$$

or more formally

$$Af(x) := \int_{\mathbb{R}^n} a(x, \xi) \hat{f}(\xi) e^{i\langle x, \xi \rangle} d\xi.$$

Such operators are called **pseudodifferential operators**. A pseudodifferential operator $A = \text{Op}(a)$ on \mathbb{R}^n is defined by its symbol $a(x, \xi)$, where a is a suitable function defined on $U \times \mathbb{R}^n \subseteq \mathbb{R}^n \times \mathbb{R}^n$ for some open set U . Here $U \times \mathbb{R}^n$ should be interpreted as the cotangent bundle T^*U . If the function a has an asymptotic expansion¹

$$a(x, \xi) \sim \sum_{i=-\infty}^m a_i(x, \xi)$$

¹Here the asymptotic expansion should be interpreted as: for every k , outside some compact set, the remainder $a - \sum_{i=k}^m a_i$ has order lower than k .

for some integer m , where each a_i is homogeneous of order i in ξ :

$$a_i(x, \lambda\xi) = \lambda^i a_i(x, \xi),$$

then $A = \text{Op}(a)$ is called a **classical pseudodifferential operator of order m** .²

The **principal symbol** of an order- m classical pseudodifferential operator is the “leading term” of the symbol:

$$\sigma(A)(x, \xi) := a_i(x, \xi) = \lim_{\lambda \rightarrow \infty} a(x, \lambda\xi)/\lambda^m$$

The constructions above can be generalised to manifolds by requiring them to be (pseudo)differential operators in local charts.³ Then (x, ξ) are local coordinates of points in T^*M , and the principal symbol $\sigma(A)$ defines a (smooth) function $T^*M \rightarrow \mathbb{C}$. We restrict it to the cosphere bundle $S^*M = \{x \in T^*M \mid \|x\| = 1\}$ to obtain $\sigma(A) \in C^\infty(S^*M)$.

If we restrict to classical pseudodifferential operators of order 0, then they are bounded operators on $L^2(M)$, and we take the norm completion to obtain a C^* -algebra $\Psi^0(M)$. Operators in $\Psi^0(M)$ are called **pseudodifferential operators of order 0**. The principal symbol σ extends to a **surjective** $*$ -homomorphism $\Psi^0(M) \rightarrow C(S^*M)$. In particular, we have the following extension of C^* -algebras, referred to as the pseudodifferential extension:



Theorem 2.4

There is an extension of C^* -algebras

$$\mathbb{K}(L^2(M)) \hookrightarrow \Psi^0(M) \xrightarrow{\sigma} C(S^*M).$$

In particular, it fits in the following commutative diagram:

$$\begin{array}{ccccc} \mathbb{K}(L^2(M)) & \hookrightarrow & \Psi^0(M) & \xrightarrow{\sigma} & C(S^*M) \\ \parallel & & \downarrow & & \downarrow \\ \mathbb{K}(L^2(M)) & \hookrightarrow & \mathbb{B}(L^2(M)) & \twoheadrightarrow & \mathbb{B}(L^2(M))/\mathbb{K}(L^2(M)). \end{array}$$

Therefore, we may identify a principal symbol $\sigma(A)$ in $C(S^*M)$ with an element in the Calkin algebra \mathbb{B}/\mathbb{K} . By functoriality of K-theory we have

$$\begin{array}{ccc} K_1(C(S^*M)) & \xrightarrow{\partial} & K_0(\mathbb{K}) \\ \downarrow & & \parallel \\ K_1(\mathbb{B}/\mathbb{K}) & \xrightarrow{\partial} & K_0(\mathbb{K}) \end{array} \quad (1)$$

So the boundary map can be identified with the Fredholm index when acting on the K_1 -class of a Fredholm operator.

Now we introduce ellipticity:



Definition 2.5

$D \in \Psi^0(M)$ is called **elliptic**, if $\sigma(D)$ is invertible on S^*M , or equivalently if $\sigma(D)(x, \xi) \neq 0$ for

²Or we can characterise the order using the growth condition: there exists constant $C_{\alpha, \beta}$ such that $|\partial_x^\beta \partial_\xi^\alpha p(x, \xi)| \leq C_{\alpha, \beta} (1 + |\xi|)^{m - |\alpha|}$ for all multi-index α and β . The order of A is the supremum of such m .

³For differential operators, which are operators with polynomial symbols, such constructions are straightforward because all differential operators are **local**. But pseudodifferential operators are not local in general. A well-defined global pseudodifferential operator on M should satisfy that it restricts to pseudodifferential operator on \mathbb{R}^n for **any** local chart of M . See [Tre13].

$(x, \xi) \in T^*M \setminus 0$.

Now Atkinson's theorem immediately implies:



Theorem 2.6

If $D \in \Psi^0(M)$ is elliptic, then it is Fredholm (as a bounded operator on $L^2(M)$). The Fredholm index $\text{Index}(D)$ of D is the image of $[\sigma(D)] \in K_1(C(S^*M))$ under the boundary map $K_1(C(S^*M)) \xrightarrow{\partial} K_0(\mathbb{K})$ in (1).

2.3 The $K(TM)$ -symbol class

We have seen that the symbol $\sigma(D)$ represents a class $[\sigma(D)] \in K_1(C(S^*M)) = K^{-1}(S^*M)$. In most literatures and in the construction of the topological index in Section 3, we consider a class in $K^0(TM)$. Given $[\sigma(D)] \in K_1(C(S^*M))$, we obtain a class in $K_0(C_0(TM)) = K(TM)$, which we also denote by $[\sigma(D)]$, under the following composition of maps:

$$K^{-1}(S^*M) \xrightarrow{\partial} K^0(T^*M) \xrightarrow{\cong} K^0(TM),$$

where:

- ∂ is the boundary map in K-theory induced by the extension

$$C_0(T^*M) \hookrightarrow C(D^*M) \rightarrow C(S^*M) \quad (2)$$

where $D^*M := \{x \in T^*M \mid \|x\| \leq 1\}$ is the **codisk bundle** over M , and $S^*M := \{x \in T^*M \mid \|x\| = 1\}$ is the **cosphere bundle** over M . The map $C(D^*M) \rightarrow C(S^*M)$ is given by restriction of functions, and the kernel consists of functions on D^*M vanishing on the boundary. D^*M is homeomorphic to the one-point compactification of T^*M , and S^*M is sent to infinity under this homeomorphism. Hence we may identify the kernel with functions on T^*M vanishing at infinity.

- We use a Riemannian metric to identify T^*M with TM , hence $K(T^*M)$ with $K(TM)$.

We have defined the analytical index $\text{a-ind}(D)$ using the class $[\sigma(D)] \in K^{-1}(S^*M)$. It turns out that it does not matter if we move to $[\sigma(D)] \in K(TM)$. In fact, we have:



Proposition 2.7


$\text{a-ind}(D)$ depends only on the $K(TM)$ -symbol class $[\sigma(D)] \in K(TM)$.



Proof (sketch)

The extension (2) induces a long exact sequence

$$\cdots \rightarrow K_1(C(D^*M)) \rightarrow K_1(C(S^*M)) \xrightarrow{\partial} K_0(C_0(T^*M)) \rightarrow \cdots,$$

if $[\sigma(D)] \in K_1(C(S^*M))$ lies in the kernel of the boundary map $K_1(C(S^*M)) \xrightarrow{\partial} K_0(C_0(T^*M))$, by exactness it is the restriction of an invertible function in $C(D^*M)$. But D^*M has properly contractible fibres, hence any function in $C(D^*M)$ is properly homotopic to a function which is fibrewise constant. Such a function is the symbol of a nowhere vanishing multiplication operator, and its index is zero since it is invertible. 

From now on, we will only consider $[\sigma(D)] \in K(TM)$. We call it the $K(TM)$ -symbol class of D .

3 The topological index

Let $[\sigma(D)]$ be the $K(TM)$ -symbol class of D . The topological index $t\text{-ind}(D)$ is defined as the image of $[\sigma(D)]$ under the following composition of maps:

$$K(TM) \xrightarrow{i_!} K(T\mathbb{R}^n) \xrightarrow{\cong} \mathbb{Z},$$

where the map $i_!$ is the **wrong-way** map, constructed using the Thom isomorphism, which we will elaborate on. The isomorphism $K(T\mathbb{R}^n) \xrightarrow{\cong} \mathbb{Z}$ is the **Bott periodicity**. In fact, it is also a special case of the Thom isomorphism.

3.1 Thom isomorphism

Let $E \rightarrow M$ be an oriented vector bundle over a compact base. The Thom isomorphism of compactly-supported cohomology states that the cohomology groups of E are isomorphic to those of M with a dimension shift.

Being more precise. Recall that a real vector bundle $E \xrightarrow{\pi} M$ of rank- k is oriented, if and only if there exists $\psi_E \in H_c^k(E)$ such that ψ_E restricts to a generator of $H_c^k(E_x)$ for each $x \in M$. Such ψ_E is unique up to a sign and called the **Thom class** of E . There is an isomorphism

$$H_c^\bullet(M) \xrightarrow{\cong} H_c^{\bullet+k}(E), \quad \alpha \mapsto \pi^* \alpha \smile \psi_E.$$

The situation in (compactly-supported) K-theory is just similar. In fact, this can be defined for all generalised cohomology theories. We say a real vector bundle $E \xrightarrow{\pi} M$ of **even** rank is **K-oriented**, if there exists $\phi_E \in K(E)$, such that ϕ_E restricts to a generator of each $K(E_x) \cong \mathbb{Z}$ (here we need that E_x is even-dimensional). There is an isomorphism (here we apply Bott periodicity to remove the dimension shift):

$$K(M) \xrightarrow{\cong} K(E), \quad \alpha \mapsto \pi^* \alpha \smile \phi_E.$$

Real vector bundles are not K-oriented in general. But we have the following:



Proposition 3.1

Every complex vector bundle is oriented and K-oriented.



Example 3.2

Bott periodicity is a special case of the Thom isomorphism. Bott periodicity states that $K(\text{pt}) \cong K(\mathbb{R}^{2n})$. Since $\mathbb{R}^{2n} \cong \mathbb{C}^n$, it is a K-oriented vector bundle over the base, which is a single point. The Thom class is just a generator $\phi_{\mathbb{R}^{2n}} \in K(\mathbb{R}^{2n})$, called the **Bott class**, and the Thom isomorphism sends an element $\alpha \in K(\text{pt})$ to $\pi^* \alpha \smile \phi_{\mathbb{R}^{2n}}$. If we identify $K(\text{pt})$ with \mathbb{Z} , then the isomorphism is just given by $n \mapsto n\phi_{\mathbb{R}^{2n}}$ for $n \in \mathbb{Z}$.

Now we would like to apply the Thom isomorphism to obtain a wrong-way map in K-theory. Let $i: X \rightarrow Y$ be a proper embedding of manifolds. Then it induces an embedding $i_*: TX \rightarrow TY$. We want to use Thom isomorphism to find a map $K(TX) \rightarrow K(TY)$ between K-theory. But TY is not a vector bundle over TX . For this we consider the extension of vector bundles over X :

$$TX \xrightarrow{i_*} TY|_X \rightarrow TY|_X/TX.$$

The vector bundle $TY|_X/TX$ is called the **normal bundle** of X in Y . We denote it by NX . Then $TY|_X$ decomposes as $TX \oplus NX$. Using the lemma,

**Lemma 3.3 (Tubular neighbourhood theorem)**

There is an open neighbourhood N of X in Y which is diffeomorphic to NX . Such an N is called a tubular neighbourhood of X in Y ,

we can identify N with the vector bundle NX over X , and N is an open subset of Y . The embedding i decomposes as $i: X \hookrightarrow N \hookrightarrow Y$, similarly $i_*: TX \hookrightarrow TN \hookrightarrow TY$. N being a vector bundle over X implies that TN is a vector bundle over TX . In particular, TN admits a complex structure and hence K -oriented: recall that $TY|_X = TX \oplus NX \cong TX \oplus N$. So $TTY|_{TX} \cong TTX \oplus TN$. But $TTY|_{TX} \cong \pi^*(TY|_X) \oplus \pi^*(TY|_X)$ and $TTX \cong \pi^*TX \oplus \pi^*TX$, where $\pi: TX \rightarrow X$ is the projection of the tangent bundle. So $TN \cong \pi^*N \oplus \pi^*N = \pi^*N \otimes \mathbb{C}$. Therefore TN can be equipped with a complex structure and we obtain the Thom isomorphism

$$K(TX) \xrightarrow{\cong} K(TN).$$

The inclusion $TN \hookrightarrow TY$ induces a map

$$K(TN) \rightarrow K(TY).$$

To see this, notice that there is an “extension by zero” $*$ -homomorphism $C_0(TN) \hookrightarrow C_0(TY)$ (here $TN \subseteq TY$ must be open!) Then we have an induced map $K_0(C_0(TN)) \rightarrow K_0(C_0(TY))$ because K -theory of C^* -algebras is functorial.

The composition of the two maps above gives a map $i_!: K(TX) \rightarrow K(TY)$ as desired.

3.2 The cohomological formula

Now we derive the cohomological formula for the topological index. This is done by a diagram chase, as first carried out in [AS3]. Recall from Bram’s talk that the diagram

$$\begin{array}{ccc} K(M) & \xrightarrow{\text{Thom}} & K(E) \\ \text{Ch} \downarrow & \color{red}{\times} & \downarrow \text{Ch} \\ H_c^{\text{even}}(M) & \xrightarrow{\text{Thom}} & H_c^{\text{even}}(E) \end{array}$$

fails to commute. The “defect” measured by the **Todd class** of \bar{E} , the conjugate bundle of E :

**Lemma 3.4**

The following diagram commutes:

$$\begin{array}{ccc} K(M) & \xrightarrow{\text{Thom}} & K(E) \\ (-1)^k \text{Td}(\bar{E})^{-1} \text{Ch} \downarrow & & \downarrow \text{Ch} \\ H_c^{\text{even}}(M) & \xrightarrow{\text{Thom}} & H_c^{\text{even}}(E) \end{array}$$

where \bar{E} is the conjugate bundle of E , and k is the **complex rank** of E .

The following diagram commutes:

$$\begin{array}{ccccccc} [\sigma(D)] & & & & \text{t-ind}(D) \cdot \phi_{\mathbb{TR}^n} & & \text{t-ind}(D) \\ K(TM) & \xrightarrow{\text{Thom}} & K(TN) & \xrightarrow{\quad} & K(\mathbb{TR}^n) & \xleftarrow{\text{Thom}} & K(\text{pt}) \cong \mathbb{Z} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_c^{\text{even}}(TM) & \xrightarrow{\text{Thom}} & H_c^{\text{even}}(TN) & \xrightarrow{\quad} & H_c^{\text{even}}(\mathbb{TR}^n) & \xleftarrow{\text{Thom}} & H_c^{\text{even}}(\text{pt}) \cong \mathbb{Q} \end{array}$$

$$(-1)^{n-m} \pi^* \text{Td}(TM \otimes \mathbb{C}) \text{Ch}([\sigma(D)]) \quad (-1)^{n-m} \pi^* \text{Td}(TM \otimes \mathbb{C}) \text{Ch}([\sigma(D)]) \text{Ch}(\psi_{TN}) = (-1)^n \text{t-ind}(D) \psi_{\mathbb{TR}^n} \quad (-1)^n \text{t-ind}(D)$$

- Let M be an m -dimensional compact manifold, embedded in \mathbb{R}^n . We have seen that N is isomorphic to the normal bundle of X in Y , and TN has complex rank $(n - m)$.
- The topological index is the image of $[\sigma(D)]$ under the composition of maps in the first row. In particular, the Thom isomorphism $K(\text{pt}) \cong K(\mathbb{R}^n)$ sends the index $\text{t-ind}(D)$ to $\text{t-ind}(D) \cdot \phi_{\mathbb{R}^n}$ where $\phi_{\mathbb{R}^n}$ is the Bott element of \mathbb{R}^n .
- We apply Chern characters to every K-theory group, and use the previous theorem to obtain this big commutative diagram. In the commutative square on the right, the Todd class of \mathbb{R}^n is just 1, and the image in $H_c^{\text{even}}(\mathbb{R}^n)$ is just $(-1)^n \text{t-ind}(D) \psi_{\mathbb{R}^n}$, where $\psi_{\mathbb{R}^n}$ is the Bott class in $H_c^{2n}(\mathbb{R}^{2n})$. The extra factor $(-1)^n$ comes from the Chern character defect. The diagram on the left can be worked out in a similar way.
- Now we look at the Todd class $\text{Td}(\overline{TN})$. We have seen that $TN \cong \pi^*N \otimes \mathbb{C}$, so TN is isomorphic to its complex conjugate. Since $TN \oplus TTM \cong \mathbb{R}^n$ is trivial, computation shows that $\text{Td}(TN) = \text{Td}(TTM)^{-1}$. And $TTM = \pi^*TM \otimes \mathbb{C}$. Therefore we obtain the characteristic class in $H_c^{\text{even}}(TN)$:

$$(-1)^{n-m} \pi^* \text{Td}(TM \otimes \mathbb{C}) \smile \text{Ch}([\sigma(D)]) \smile \psi_{TN},$$

whose image in $H_c^{\text{even}}(\mathbb{R}^n)$ is just $\text{t-ind}(D) \cdot \phi_{\mathbb{R}^n}$ since the diagram commutes.

- The last thing is to unzip t-ind from the characteristic classes we have just obtained. We integrate the two forms over \mathbb{R}^n (which is just pairing with the fundamental class of \mathbb{R}^n in $H_{2n}(\mathbb{R}^n)$):

$$\int_{\mathbb{R}^n} (-1)^{n-m} \pi^* \text{Td}(TM \otimes \mathbb{C}) \smile \text{Ch}([\sigma(D)]) \smile \psi_{TN} = \int_{\mathbb{R}^n} (-1)^n \text{t-ind}(D) \psi_{\mathbb{R}^n}.$$

The right-hand side is just $(-1)^n \text{t-ind}(D)$. The integral on the left-hand side restricts to the integral on TN since ψ_{TN} is compactly supported on TN . Recall that the inverse of the Thom isomorphism is integrating along the fibres, the integral becomes

$$(-1)^{n-m} \int_{TM} \pi^* \text{Td}(TM \otimes \mathbb{C}) \smile \text{Ch}([\sigma(D)]).$$

Compare both sides of the equation we obtain:



Theorem 3.5 (Cohomological formula of the topological index, I)

The topological index $\text{t-ind}(D)$ equals

$$\text{t-ind}(D) = (-1)^m \int_{TM} \pi^* \text{Td}(TM \otimes \mathbb{C}) \smile \text{Ch}([\sigma(D)]),$$

where m is the dimension of M , $\pi: TM \rightarrow M$ is the bundle projection of TM .

If M is oriented, then Thom isomorphism applies to TM and we can obtain a similar cohomological formula where the integral is on M instead of TM . However, an extra factor $(-1)^{m(m-1)/2}$ is involved due to the orientation of M .



Theorem 3.6 (Cohomological formula of the topological index, II)

If M is oriented, then the topological index $\text{t-ind}(D)$ equals

$$\text{t-ind}(D) = (-1)^{m(m+1)/2} \int_M \pi_!(\pi^* \text{Td}(TM \otimes \mathbb{C}) \smile \text{Ch}([\sigma(D)])),$$

where $\pi_!$ is “integrating along fibres”.

4 Remarks on the general cases

So far, we have imposed several restrictions on the pseudodifferential operator D .

- The **ellipticity** guarantees that D is Fredholm. This is indeed not a necessary condition: there are operators which are not elliptic but nevertheless Fredholm, see for example [BvE14]. This is, however, far from our goal.
- We have restricted to **order-zero** operators. Such operators are bounded on $L^2(M)$. While considering pseudodifferential operators of higher orders, they are no longer bounded on $L^2(M)$. An elliptic pseudodifferential operator D of order m defines a bounded Fredholm operator $H^s(M) \rightarrow H^{s-m}(M)$ for all s , where $H^s(M)$ and $H^{s-m}(M)$ are Sobolev spaces. Another viewpoint is to consider D as an **unbounded** Fredholm operator on $L^2(M)$.

In either case, a pseudodifferential operator D of order greater than 0 does not belong to $\mathbb{B}(L^2(M))$; such operators do not even form an algebra. But we can consider the operator

$$D(1 + D^2)^{-1/2},$$

which has order zero, hence bounded. In particular, this operator has the same symbol class with D . Since indices are defined as collections of maps $K(TX) \rightarrow \mathbb{Z}$, we see that studying order-zero operators is enough.

- We have restricted to **scalar-valued** operators, that is, they are defined by scalar-valued symbols. However, most elliptic differential operators are operators which act on sections of vector bundles; that is, they are linear maps

$$D: C^\infty(M; E) \rightarrow C^\infty(M; E)$$

where E is a vector bundle over a compact manifold M , and $C^\infty(M; E)$ denotes the smooth sections. (The scalar valued case can be viewed as such a map with $E = M \times \mathbb{C}$). In these cases, the symbol $\sigma(D)$ is a section of the endomorphism bundle

$$\text{End}(\pi^* E),$$

where $\pi: T^*M \rightarrow M$ is the bundle projection; or rather its restriction to S^*M . Such sections are in bijection with

$$\text{End}_{C(S^*M)} C(S^*M; \pi^* E),$$

the **C*-algebra of adjointable operators** on the **Hilbert C(S*M)-module** $C(S^*M; \pi^* E)$. Assuming M is compact, then S^*M is also compact and hence $C(S^*M)$ is unital. Since $\pi^* E$ is finite-dimensional, $C(S^*M; \pi^* E)$ is finitely-generated. Then $\text{End}_{C(S^*M)} C(S^*M; \pi^* E)$ is Morita equivalent to $C(S^*M)$. Consequently, nothing changes on the K-theory level. See, e.g. [MT05], for more on Hilbert C*-modules.

5 Proof of the index theorem

In the end, we will sketch the elegant proof of the index theorem, due to Atiyah–Singer [AS1]. The strategy of the proof is to show that there is a unique index satisfying certain constraints. Both a-ind and t-ind satisfy them, hence the two indices must coincide.

More precisely, an index is a collection of group homomorphisms $\text{ind} = \{K(TX) \rightarrow \mathbb{Z}\}$ for each compact manifold X . We have



Proposition 5.1

There is a unique index ind satisfying 1 and 2:

1. If $X = \text{pt}$, then $\text{ind}: K(\text{pt}) \rightarrow \mathbb{Z}$ is the identity map.
2. If $X \hookrightarrow Y$ is a proper embedding of manifolds, then $\text{ind} \circ i_! = \text{ind}$, where $i_!$ is the wrong-way map. (See Section 3.1).



Proof

Consider $X \xrightarrow{i} \mathbb{R}^n \xleftarrow{j} \text{pt}$, where $X \hookrightarrow \mathbb{R}^n$ is a proper embedding. It induces

$$\begin{array}{ccccc}
 K(TX) & \xrightarrow{i_!} & K(T\mathbb{R}^n) & \xleftarrow{j_!} & K(\text{pt}) \\
 & \searrow \text{ind} & \downarrow \text{ind} & \swarrow \text{ind} & \\
 & & \mathbb{Z} & &
 \end{array} ,$$

Condition 2 claims that the index $K(TX) \rightarrow \mathbb{Z}$ on X is determined by the index $K(T\mathbb{R}^n) \rightarrow \mathbb{Z}$ on \mathbb{R}^n , and by Thom isomorphism determined by the index $K(\text{pt}) \rightarrow \mathbb{Z}$ on pt , but this is unique by Condition 1.

The remaining work is to show that both a-ind and t-ind satisfy Conditions 1–2 in the previous proposition.



Proposition 5.2

t-ind satisfies conditions 1–2.



Proof

1 is just Bott periodicity; 2 is by functoriality.



Proposition 5.3

a-ind satisfies conditions 1–2.

Condition 1 for a-ind is easy to check. Let $M = \text{pt}$, then a vector bundle over M is just a finite-dimensional vector space, and a class $[E] - [F] \in K(\text{pt})$ is identified with $\dim E - \dim F \in \mathbb{Z}$. A symbol $\sigma(D)$ is just a linear map $E \rightarrow F$, so $\dim E = \dim \ker D + \dim \text{im } D = \dim \ker D + \dim F - \dim \text{coker } D$. Hence $\text{a-ind}(D) = \dim E - \dim F$.

Checking condition 2 is difficult. For a detailed proof we refer to Atiyah–Singer’s original work [AS1], in which they introduced the equivariant index theorem as well. Recall that $i_!$ is constructed as the composition

$$\begin{array}{ccccc}
 i_!: K(TX) & \xrightarrow[\cong]{\text{Thom}} & K(TN) & \longrightarrow & K(TY) \\
 & \searrow \text{ind} & \downarrow \text{ind} & \swarrow \text{ind} & \\
 & & \mathbb{Z} & &
 \end{array}$$

To show that the outer diagram commutes, it suffices to show that the two triangles commute. The right triangle commutes due to the **excision** property of the analytical index. The idea: the

extension homomorphism $C_0(TU) \rightarrow C_0(TX)$ lifts to $C(S^*U) \rightarrow C(S^*X)$, and the boundary map in K-theory is natural, so we must have the same index.

The commutativity of the left triangle is hard and we will only sketch the main ideas. The proof is based on the following “product formula”, and its equivariant version.

Recall that K-theory of topological spaces allows an exterior product (cross product):

$$K(X) \otimes K(Y) \xrightarrow{\times} K(X \times Y), \quad x \otimes y \mapsto \pi_X^* x \smile \pi_Y^* y$$

where $\pi_X: X \times Y \rightarrow X$ and $\pi_Y: X \times Y \rightarrow Y$.



Lemma 5.4

Let $x \in K(TX)$ and $y \in K(TY)$. The the cross product $x \times y$ is an element in $K(TX \times TY) \cong K(T(X \times Y))$. And we have

$$\text{a-ind}(x \times y) = \text{a-ind}(x) \cdot \text{a-ind}(y).$$

The product formula already solves the case when N is **trivial**: the Thom isomorphism

$$K(TX) \rightarrow K(TN)$$

sends $x \in K(TX)$ to $\pi_1^* x \cdot \phi_{TN}$ where π_1 is the projection $TN \rightarrow TX$. If $N \cong X \times \mathbb{R}^n$ is trivial, then ϕ_{TN} is just the pullback of the Bott element $\phi_{\mathbb{R}^n} \in K(\mathbb{R}^n)$ under the projection $TN \xrightarrow{p} \mathbb{R}^n$. So x is sent to $\pi_1^* x \smile \pi_2^* \phi_{\mathbb{R}^n} = x \times \phi_{\mathbb{R}^n}$. The product formula then claims that $\text{a-ind}(x \times \phi_{\mathbb{R}^n}) = \text{a-ind}(x) \cdot \text{a-ind}(\phi_{\mathbb{R}^n}) = \text{a-ind}(x)$ because the Bott element has index 1.

The discussion above also shows that the Thom isomorphism for the trivial bundle $TN \cong TX \times \mathbb{R}^n \cong TX \times \mathbb{R}^{2n}$ is given by the composition

$$K(TX) \rightarrow K(TX) \otimes K(\mathbb{R}^{2n}) \xrightarrow{\times} K(TN), \quad x \mapsto x \otimes \phi_{\mathbb{R}^{2n}} \mapsto x \times \phi_{\mathbb{R}^{2n}}.$$

But N is not trivial in general. In that case, the cross product $K(TX) \otimes K(\mathbb{R}^n) \xrightarrow{\times} K(TN)$ does not make sense. But one can define a “twisted” cross product instead. In this situation we require the K-theory classes in the typical fibre \mathbb{R}^n are well-defined globally, this requires they are **equivariant** with respect to the structure group $O(n)$. The cross product should be replaced by a twisted product

$$K(TX) \otimes K_{O(n)}(\mathbb{R}^n) \rightarrow K(TN)$$

where we need an “ $O(n)$ -equivariant Bott element” $\phi_{\mathbb{R}^n}^{O(n)}$, which comes from the following “equivariant Bott periodicity”:

$$K_{O(n)}(\text{pt}) \xrightarrow{\cong} K_{O(n)}(\mathbb{R}^n), \quad x \mapsto \pi^* x \smile \phi_{\mathbb{R}^n}^{O(n)}.$$

Atiyah and Singer proved that an “equivariant product formula” holds as well, and the Thom isomorphism $K(TX) \xrightarrow{\cong} K(TN)$ sends an element to its twisted product with the equivariant Bott element. Since the equivariant Bott element has analytical index 1, a similar argument as in the trivial case shows that the Thom isomorphism does not change the analytical index, and the proof is done.

Notes and comments

I assume that my audience have some basic knowledge on K-theory and algebraic topology. Knowing K-theory for C^* -algebras will be helpful but not necessary, as we will only use very few properties of them.

Most of the contents and details can be found in Atiyah–Singer’s masterpiece [AS1]. The cohomological formula is presented in [AS3]. Literature that I refer to include [Muk13, Sha06, Lan05], but they are mostly explaining the original article by Atiyah and Singer.

The thesis [vE00] should be especially mentioned, since it interprets the index theory in the framework of K-theory of C^* -algebras, which I favour. Such a strategy should have a long history, and eventually leads to the most elegant and powerful way to attack the index theory: Kasparov’s bivariant K-theory [Kas88]. I have to admit that although K-theory of C^* -algebras is mentioned in my talk, the proofs depend quite little on this powerful tool. One can find a purely C^* -algebraic proof in [Hig93], which is closely related to E-theory, another bivariant K-theory of C^* -algebras.

References

- [AS1] Michael F Atiyah and Isadore Manuel Singer. The index of elliptic operators: I. *Annals of mathematics*, pages 484–530, 1968.
- [AS3] Michael F Atiyah and Isadore M Singer. The index of elliptic operators: III. *Annals of mathematics*, pages 546–604, 1968.
- [AS4] Michael F Atiyah and Isadore M Singer. The index of elliptic operators: IV. *Annals of Mathematics*, 93(1):119–138, 1971.
- [BvE14] Paul F Baum and Erik van Erp. K-homology and index theory on contact manifolds. *Acta Mathematica*, 213(1):1–48, 2014.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge: Cambridge University Press, 2002.
- [Hat03] Allen Hatcher. *Vector bundles and K-theory*. <http://www.math.cornell.edu/~hatcher>, 2003.
- [Hig93] Nigel Higson. On the K-theory proof of the index theorem. *Contemporary Mathematics*, 148:67–67, 1993.
- [Kas88] Gennadi G Kasparov. Equivariant KK-theory and the Novikov conjecture. *Inventiones mathematicae*, 91(1):147–201, 1988.
- [Lan05] Gregory D Landweber. K-theory and elliptic operators. *arXiv preprint math/0504555*, 2005.
- [MT05] Vladimir M Manuilov and Evgenij V Troitsky. Hilbert C^* -modules, translations of mathematical monographs, 226. *American Mathematical Society, Providence, RI*, 2005.
- [Muk13] Amiya Mukherjee. *Atiyah–Singer Index Theorem – An Introduction*. Springer, 2013.
- [Ols94] Niels Erik Wegge Olsen. *K-theory and C^* -algebras: A Friendly Approach*. Oxford University Press, 1994.
- [Sha06] Patrick Shanahan. *The Atiyah–Singer index theorem: an introduction*, volume 638. Springer, 2006.
- [Tre13] Jean-François Treves. *Introduction to Pseudodifferential and Fourier Integral Operators: Pseudodifferential Operators*. Springer Science & Business Media, 2013.
- [vE00] Erik van Erp. The Atiyah–Singer index theorem, C^* -algebraic K-theory, and quantization. 2000. Master thesis.