

In Exercises 1–6, determine which sets of vectors are orthogonal.

1.  $\begin{bmatrix} -1 \\ 4 \\ -3 \end{bmatrix}, \begin{bmatrix} 5 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ -4 \\ -7 \end{bmatrix}$       2.  $\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -5 \\ -2 \\ 1 \end{bmatrix}$

3.  $\begin{bmatrix} 2 \\ -7 \\ -1 \end{bmatrix}, \begin{bmatrix} -6 \\ -3 \\ 9 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix}$       4.  $\begin{bmatrix} 2 \\ -5 \\ -3 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ -2 \\ 6 \end{bmatrix}$

In Exercises 7–10, show that  $\{\mathbf{u}_1, \mathbf{u}_2\}$  or  $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$  is an orthogonal basis for  $\mathbb{R}^2$  or  $\mathbb{R}^3$ , respectively. Then express  $\mathbf{x}$  as a linear combination of the  $\mathbf{u}$ 's.

7.  $\mathbf{u}_1 = \begin{bmatrix} 2 \\ -3 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 6 \\ 4 \end{bmatrix},$  and  $\mathbf{x} = \begin{bmatrix} 9 \\ -7 \end{bmatrix}$

10.  $\mathbf{u}_1 = \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix},$  and  $\mathbf{x} = \begin{bmatrix} 5 \\ -3 \\ 1 \end{bmatrix}$

12. Compute the orthogonal projection of  $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$  onto the line through  $\begin{bmatrix} -1 \\ 3 \end{bmatrix}$  and the origin.

14. Let  $\mathbf{y} = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$  and  $\mathbf{u} = \begin{bmatrix} 7 \\ 1 \end{bmatrix}$ . Write  $\mathbf{y}$  as the sum of a vector in  $\text{Span}\{\mathbf{u}\}$  and a vector orthogonal to  $\mathbf{u}$ .
15. Let  $\mathbf{y} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$  and  $\mathbf{u} = \begin{bmatrix} 8 \\ 6 \end{bmatrix}$ . Compute the distance from  $\mathbf{y}$  to the line through  $\mathbf{u}$  and the origin.

In Exercises 17–22, determine which sets of vectors are orthonormal. If a set is only orthogonal, normalize the vectors to produce an orthonormal set.

17.  $\begin{bmatrix} 1/3 \\ 1/3 \\ 1/3 \end{bmatrix}, \begin{bmatrix} -1/2 \\ 0 \\ 1/2 \end{bmatrix}$       18.  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$

True false (plus justification):

24. a. Not every orthogonal set in  $\mathbb{R}^n$  is linearly independent.  
 b. If a set  $S = \{\mathbf{u}_1, \dots, \mathbf{u}_p\}$  has the property that  $\mathbf{u}_i \cdot \mathbf{u}_j = 0$  whenever  $i \neq j$ , then  $S$  is an orthonormal set.  
 c. If the columns of an  $m \times n$  matrix  $A$  are orthonormal, then the linear mapping  $\mathbf{x} \mapsto A\mathbf{x}$  preserves lengths.  
 d. The orthogonal projection of  $\mathbf{y}$  onto  $\mathbf{v}$  is the same as the orthogonal projection of  $\mathbf{y}$  onto  $c\mathbf{v}$  whenever  $c \neq 0$ .  
 e. An orthogonal matrix is invertible.
26. Suppose  $W$  is a subspace of  $\mathbb{R}^n$  spanned by  $n$  nonzero orthogonal vectors. Explain why  $W = \mathbb{R}^n$ .

30. Let  $U$  be an orthogonal matrix, and construct  $V$  by interchanging some of the columns of  $U$ . Explain why  $V$  is an orthogonal matrix.

In Exercises 1 and 2, you may assume that  $\{\mathbf{u}_1, \dots, \mathbf{u}_4\}$  is an orthogonal basis for  $\mathbb{R}^4$ .

1.  $\mathbf{u}_1 = \begin{bmatrix} 0 \\ 1 \\ -4 \\ -1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} 3 \\ 5 \\ 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ -4 \end{bmatrix}$ ,  $\mathbf{u}_4 = \begin{bmatrix} 5 \\ -3 \\ -1 \\ 1 \end{bmatrix}$ ,  
 $\mathbf{x} = \begin{bmatrix} 10 \\ -8 \\ 2 \\ 0 \end{bmatrix}$ . Write  $\mathbf{x}$  as the sum of two vectors, one in  $\text{Span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$  and the other in  $\text{Span}\{\mathbf{u}_4\}$ .

In Exercises 3–6, verify that  $\{\mathbf{u}_1, \mathbf{u}_2\}$  is an orthogonal set, and then find the orthogonal projection of  $\mathbf{y}$  onto  $\text{Span}\{\mathbf{u}_1, \mathbf{u}_2\}$ .

3.  $\mathbf{y} = \begin{bmatrix} -1 \\ 4 \\ 3 \end{bmatrix}$ ,  $\mathbf{u}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$

4.  $\mathbf{y} = \begin{bmatrix} 6 \\ 3 \\ -2 \end{bmatrix}$ ,  $\mathbf{u}_1 = \begin{bmatrix} 3 \\ 4 \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -4 \\ 3 \\ 0 \end{bmatrix}$

In Exercises 7–10, let  $W$  be the subspace spanned by the  $\mathbf{u}$ 's, and write  $\mathbf{y}$  as the sum of a vector in  $W$  and a vector orthogonal to  $W$ .

$$7. \mathbf{y} = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}, \mathbf{u}_1 = \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 5 \\ 1 \\ 4 \end{bmatrix}$$

In Exercises 11 and 12, find the closest point to  $\mathbf{y}$  in the subspace  $W$  spanned by  $\mathbf{v}_1$  and  $\mathbf{v}_2$ .

$$11. \mathbf{y} = \begin{bmatrix} 3 \\ 1 \\ 5 \\ 1 \end{bmatrix}, \mathbf{v}_1 = \begin{bmatrix} 3 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}$$

In Exercises 13 and 14, find the best approximation to  $\mathbf{z}$  by vectors of the form  $c_1\mathbf{v}_1 + c_2\mathbf{v}_2$ .

$$13. \mathbf{z} = \begin{bmatrix} 3 \\ -7 \\ 2 \\ 3 \end{bmatrix}, \mathbf{v}_1 = \begin{bmatrix} 2 \\ -1 \\ -3 \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ -1 \end{bmatrix}$$

15. Let  $\mathbf{y} = \begin{bmatrix} 5 \\ -9 \\ 5 \end{bmatrix}$ ,  $\mathbf{u}_1 = \begin{bmatrix} -3 \\ -5 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -3 \\ 2 \\ 1 \end{bmatrix}$ . Find the distance from  $\mathbf{y}$  to the plane in  $\mathbb{R}^3$  spanned by  $\mathbf{u}_1$  and  $\mathbf{u}_2$ .

In Exercises 21 and 22, all vectors and subspaces are in  $\mathbb{R}^n$ . Mark each statement True or False. Justify each answer.

21. a. If  $\mathbf{z}$  is orthogonal to  $\mathbf{u}_1$  and to  $\mathbf{u}_2$  and if  $W = \text{Span}\{\mathbf{u}_1, \mathbf{u}_2\}$ , then  $\mathbf{z}$  must be in  $W^\perp$ .
- b. For each  $\mathbf{y}$  and each subspace  $W$ , the vector  $\mathbf{y} - \text{proj}_W \mathbf{y}$  is orthogonal to  $W$ .
- c. The orthogonal projection  $\hat{\mathbf{y}}$  of  $\mathbf{y}$  onto a subspace  $W$  can sometimes depend on the orthogonal basis for  $W$  used to compute  $\hat{\mathbf{y}}$ .
- d. If  $\mathbf{y}$  is in a subspace  $W$ , then the orthogonal projection of  $\mathbf{y}$  onto  $W$  is  $\mathbf{y}$  itself.

In Exercises 1–6, the given set is a basis for a subspace  $W$ . Use the Gram–Schmidt process to produce an orthogonal basis for  $W$ .

$$3. \begin{bmatrix} 2 \\ -5 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 2 \end{bmatrix}$$

$$5. \begin{bmatrix} 1 \\ -4 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 7 \\ -7 \\ -4 \\ 1 \end{bmatrix}$$

7. Find an orthonormal basis of the subspace spanned by the vectors in Exercise 3.

Find an orthogonal basis for the column space of each matrix in Exercises 9–12.

9. 
$$\begin{bmatrix} 3 & -5 & 1 \\ 1 & 1 & 1 \\ -1 & 5 & -2 \\ 3 & -7 & 8 \end{bmatrix}$$

10. 
$$\begin{bmatrix} -1 & 6 & 6 \\ 3 & -8 & 3 \\ 1 & -2 & 6 \\ 1 & -4 & -3 \end{bmatrix}$$